

MATH255 - Honours Analysis 2

Basic point-set topology; metric spaces; Hölder-Minkowski Inequalities; compactness; series, series of functions, uniform and pointwise convergence.

Based on lectures from Winter, 2024 by Prof. Dimitry Jakobson

Notes by Louis Meunier

CONTENTS

- 1 INTRODUCTION** **3**
- 1.1 Metric Spaces 3
- 2 POINT-SET TOPOLOGY** **7**
- 2.1 Definitions 7
- 2.2 Basis 9
- 2.3 Subspaces 10
- 2.4 Continuous Functions 11
- 2.5 Product Spaces 12
- 2.6 Metrizable 15
- 2.7 Compactness, Connectedness 16
- 2.8 Path Components, Connected Components 21
- 2.8.1 Cantor Staircase Function 22
- 3 L^p SPACES** **23**
- 3.1 Review of ℓ^p Norms 23
- 3.2 ℓ^p Norms, Hölder-Minkowski Inequalities 23
- 3.3 Complete Metric Spaces, Completeness of ℓ_p 27
- 3.4 Contraction Mapping Theorem 29
- 3.5 Equivalent Notions of Compactness in Metric Spaces 32
- 4 DERIVATIVES** **35**
- 4.1 Introduction 35
- 4.2 Chain Rule 35
- 4.3 Critical Points 36
- 4.4 Aside: Continued Fractions 37

4.5	Back To Derivatives	38
4.6	L'Hopital's Rules	38
4.7	Taylor's Theorem	39
4.8	Convex Sets	40
5	RIEMANN INTEGRAL	40
5.1	Introduction	40
5.2	Cauchy Criterion	41
5.3	Squeeze Theorem	42
5.4	Fundamental Theorem of Calculus	44
5.5	Upper and Lower Riemann Sums	45
5.6	Indefinite Integral	45
5.7	Lebesgue Integrability Criterion	46
5.8	Integration by Parts	47
6	FUNCTION SEQUENCES, SERIES	48
6.1	Pointwise and Uniform Convergence	48
6.2	Series	50
6.3	Tests for Absolute Convergence	50
6.4	Tests for Non-Absolute Convergence	51
6.5	Series of Functions	52
6.6	Power Series	53
7	APPENDIX	55
7.1	Notes from Tutorials	55
7.2	Miscellaneous	58
7.3	Class Midterm Solutions	59
8	LIST OF THEOREMS	63

1 INTRODUCTION

1.1 Metric Spaces

↪ Definition 1.1: Metric Space

A set X is a *metric space* with distance d if

1. (symmetric) $d(x, y) = d(y, x) \geq 0$
2. $d(x, y) = 0 \iff x = y$
3. (triangle inequality) $d(x, y) + d(y, z) \geq d(x, z)$

Remark 1.1. If 1., 3. are satisfied but not 2., d can be called a “pseudo-distance”.

↪ Definition 1.2: Open Metric Space

Let (X, d) be a metric space. A subset $A \subseteq X$ is open $\iff \forall x \in A, \exists r = r(x) > 0$ s.t. $B(x, r(x)) \subseteq A$.

↪ Definition 1.3: Normed Space

Let X be a vector space over \mathbb{R} . The norm on X , denoted $\|x\| \in \mathbb{R}$, is a function that satisfies

1. $\|x\| \geq 0$
2. $\|x\| = 0 \iff x = 0$
3. $\|c \cdot x\| = |c| \cdot \|x\|$
4. $\|x + y\| \leq \|x\| + \|y\|$

If X is a normed vector space over \mathbb{R} , we can define a distance d on X by $d(x, y) = \|x - y\|$.

↪ Proposition 1.1

If X is a normed vector space over \mathbb{R} , a distance d on X by $d(x, y) = \|x - y\|$ makes (X, d) a metric space.

Proof. 1. $d(x, y) = \|x - y\| \geq 0$

2. $d(x, y) = 0 \iff \|x - y\| = 0 \iff x - y = 0 \iff x = y$

3. $d(x, y) + d(y, z) = \|x - y\| + \|y - z\| \geq \|(x - y) + (y - z)\| = \|x - z\| := d(x, z)$

⊗ **Example 1.1:** L^p distance in \mathbb{R}^n

Let $\bar{x} \in \mathbb{R}^n$, $x = (x_1, x_2, \dots, x_n)$. The L^p norm is defined

$$\|x\|_p := (|x_1|^p + |x_2|^p + \dots + |x_n|^p)^{\frac{1}{p}}.$$

In the case $p = 2, n = 2$, we simply have the standard Euclidean distance over \mathbb{R}^2 .

Unit Balls: consider when $\|x\|_p \leq 1$, over \mathbb{R}^2 .

- $p = 1$: $|x_1| + |x_2| \leq 1$; this forms a “diamond ball” in the plane.
- $p = 2$: $\sqrt{|x_1|^2 + |x_2|^2} \leq 1$; this forms a circle of radius 1. Clearly, this surrounds a larger area than in $p = 2$.

A natural question that follows is what happens as $p \rightarrow \infty$? Assuming $|x_1| \geq |x_2|$:

$$\begin{aligned} \|x\|_p &= (|x_1|^p + |x_2|^p)^{\frac{1}{p}} \\ &= \left[|x_1|^p \left(1 + \left| \frac{x_2}{x_1} \right|^p \right) \right]^{\frac{1}{p}} \\ &= |x_1| \left(1 + \left| \frac{x_2}{x_1} \right|^p \right)^{\frac{1}{p}} \end{aligned}$$

If $|x_1| > |x_2|$, this goes to $|x_1|$. If they are instead equal, then $\|x\|_p = |x_1| \cdot 2^{\frac{1}{p}} \rightarrow |x_1| \cdot 1$ as well. Hence, $\lim_{p \rightarrow \infty} \|x\|_p = \max\{|x_1|, |x_2|\}$. Thus, the unit ball will approach $\max\{|x_1|, |x_2|\} \leq 1$, that is, the unit square.

↔ **Proposition 1.2**

Let $x \in \mathbb{R}^n$. Then, $\|x\|_p \rightarrow \max\{|x_1|, \dots, |x_n|\}$ as $p \rightarrow \infty$.

Remark 1.2. This is an extension of the previous example to arbitrary real space; the proof follows nearly identically.

↔ **Definition 1.4: Convex Set**

Let X be a normed space, and take $x, y \in X$. The line segment from x to y is the set

$$\{t \cdot x + (1 - t) \cdot y : 0 \leq t \leq 1\}.$$

Let $A \subseteq X$. A is *convex* $\iff \forall x, y \in A$, we have that

$$(t \cdot x + (1 - t) \cdot y) \in A \forall 0 \leq t \leq 1.$$




Figure 1: Regions of \mathbb{R}^2 where $\|x\|_p \leq 1$ for various values of p .




Figure 2: Convex (left) versus not convex (right) sets.

Remark 1.3. Think of this as saying “a set is convex iff every point on a line segment connected any two points is in the set”.

↪ **Definition 1.5:** ℓ_p

The space ℓ_p of sequences is defined as

$$\{x = (x_1, x_2, \dots, x_n, \dots) : \sum_{n=1}^{\infty} |x_n|^p < +\infty\} \quad * .$$

Then, $*$ defines the ℓ^p norm on the space of sequences; that is, $\|x\|_p := (\sum_{n=1}^{\infty} |x_n|^p)^{\frac{1}{p}}$.

⊗ **Example 1.2:** $\ell_p, x_n = \frac{1}{n}$

. Let $x_n = \frac{1}{n}$. For which p is $x \in \ell_p$? We have, raising the norm to the power of p for ease:

$$\begin{aligned} \|x\|_p^p &= |x_1|^p + |x_2|^p + \cdots + |x_n|^p + \cdots \\ &= 1^p + \left(\frac{1}{2}\right)^p + \cdots < \infty \iff p > 1. \end{aligned}$$

In the case that $p = 1$, this becomes a harmonic sum, which diverges.

⊗ **Example 1.3:** L^p space of functions

Let $f(x)$ be a continuous function. We define the norm of f over an interval $[a, b]$

$$\|f\|_p = \left[\int_a^b |f(x)|^p dx \right]^{\frac{1}{p}}.$$

Remark 1.4. Triangle inequality for $\|x\|_p$ or $\|f\|_p$ is called Minkowski inequality; $\|x\|_p + \|y\|_p \geq \|x + y\|_p$. This will be discussed further.

⊗ **Example 1.4:** Distances between sets in \mathbb{R}^2

Let A, B be bounded, closed, “nice” sets in \mathbb{R}^2 . We define

$$d(A, B) := \text{Area}(A \Delta B),$$

where

$$A \Delta B : (A \setminus B) \cup (B \setminus A) = (A \cup B) \setminus (A \cap B).$$

It can be shown that this is a “valid” distance.

Remark 1.5. Δ denotes the “symmetric difference” of two sets.

⊗ **Example 1.5:** p -adic distance

Let p be a prime number. Let $x = \frac{a}{b} \in \mathbb{Q}$, and write $x = p^k \cdot \left(\frac{c}{d}\right)$, where c, d are not divisible by p . Then, the p -adic norm is defined $\|x\|_p := p^{-k}$. It can be shown that this is a norm.

Suppose $p = 2, x = 28 = 4 \cdot 7 = 2^2 \cdot 7$. Then, $\|28\|_2 = 2^{-2} = \frac{1}{4}$; similarly, $\|1024\|_2 = \|2^{10}\|_2 = 2^{-10}$.

More generally, we have that $\|2^k\|_2 = 2^{-k}$; conversely, $\|2^{-k}\|_2 = 2^k$. That is, the closer to 0, the larger the distance, and vice versa, contrary to our notion of Euclidean distance.

↪ Proposition 1.3

$\|x\|_p$ as defined above is a well-defined norm over \mathbb{Q} .

Proof. Left as a (homework) exercise. ■

2 POINT-SET TOPOLOGY

2.1 Definitions

↪ Definition 2.1: Topological space

A set X is a topological space if we have a collection of subsets τ of X called *open sets* s.t.

1. $\emptyset \in \tau, X \in \tau$
2. Consider $\{A_\alpha\}_{\alpha \in I}$ where A_α an open set for any α ; then, $\bigcup_{\alpha \in I} A_\alpha \in \tau$, that is, it is also an open set.
3. If J is a finite set, and A_β open for all $\beta \in J$, then $\bigcap_{\beta \in J} A_\beta \in \tau$ is also open.

In other words, 2.: arbitrary unions of open sets are open, and 3.: finite intersections of open sets are open.

↪ Definition 2.2: Closed sets

Closed sets are complements of open sets; hence, axioms for closed sets follow appropriately;

- 1.* X, \emptyset closed;
- 2.* B_α closed $\forall \alpha \in I \implies \bigcap_{\alpha \in I} B_\alpha$ closed.
- 3.* B_β closed $\forall \beta \in J, J$ finite, then $\bigcup_{\beta \in J} B_\beta$ also closed.

↪ Lecture 01; Last Updated: Tue Apr 9 14:45:17 EDT 2024

↪ **Definition 2.3: Equivalence of Metrics**

Suppose we have a metric space X with two distances d_1, d_2 ; will these necessarily admit the same topology?

A sufficient condition is that, if $\forall x \neq y \in X, \exists 1 < C < +\infty$ s.t.

$$\frac{1}{C} < \frac{d_1(x, y)}{d_2(x, y)} < C.$$

That is, the distances are equivalent, up to multiplication by a constant.

Indeed, this condition gives that $d_2 < C d_1$ and $d_2 > \frac{d_1}{C}$; this gives

$$B_{d_1}\left(x, \frac{r}{C}\right) \subseteq B_{d_2}(x, r) \subseteq B_{d_1}(x, C \cdot r).$$

Hence, d_1, d_2 define the same open/closed sets on X thus admitting the same topologies. We write $d_1 \asymp d_2$.

Remark 2.1. If $d_1 \asymp d_2$ and $d_2 \asymp d_3$, then also $d_1 \asymp d_3$. Moreover, clearly, $d_1 \asymp d_1$ and $d_1 \asymp d_2 \implies d_2 \asymp d_1$, hence this is a well-defined equivalence relation.

Hence, it's enough to show that $\forall 1 < p < +\infty$, we have $\|x\|_p \asymp \|x\|_\infty$ to show that any $\|x\|_q$ norm are equivalent for all q on \mathbb{R}^n .

↪ **Definition 2.4: Interior, Boundary of a Topological Set**

Let X be a topological space, $A \subseteq X$ and let $x \in X$. We have the following possibilities

1. $\exists U$ -open : $x \in U \subseteq A$. In this case, we say $x \in$ the *interior* of A , denoted

$$x \in \text{Int}(A).$$

2. $\exists V$ -open : $x \in V \subseteq X \setminus A = A^C$. In this case, we write

$$x \in \text{Int}(A^C).$$

3. $\forall U$ -open : $x \in U, U \cap A \neq \emptyset$ AND $U \cap A^C \neq \emptyset$. In this case, we say x is in the *boundary* of A , and denote

$$x \in \partial A.$$

↪ **Definition 2.5: Closure**

$x \in \text{Int}(A)$ or $x \in \partial A$ (that is, $x \in \text{Int}(A) \cup \partial A$) \iff every open set U that contains x intersects A .¹ Such points are called *limit points* of A . The set of all limit points of A is called the *closure* of A , denoted \bar{A} .

Remark 2.2. We have that

$$\text{Int}(A) \subseteq A \subseteq \bar{A} = \text{Int}(A) \cup \partial A.$$

↪ **Proposition 2.1: Properties of $\text{Int}(A)$**

$\text{Int}(A)$ is open, and it is the largest open set contained in A . It is the union of all U -open s.t. $U \subseteq A$. Moreover, we have that

$$\text{Int}(\text{Int}(A)) = \text{Int}(A).$$

↪ **Proposition 2.2: Properties of \bar{A}**

\bar{A} is closed; \bar{A} is the smallest closed set that contains A , that is, $\bar{A} = \bigcap B$ where B closed and $A \subseteq B$. We have too that

$$\overline{(\bar{A})} = \bar{A}.$$

↪ **Proposition 2.3**

1. A is open $\iff A = \text{Int}(A)$
2. A is closed $\iff A = \bar{A}$

2.2 Basis

↪ **Definition 2.6: Basis for a Topology**

Let τ be a topology on X . Let $\mathcal{B} \subseteq \tau$ be a collection of open sets in X such that every open set is a union of open sets in \mathcal{B} .

⊗ **Example 2.1: Example Basis**

$X = \mathbb{R}$, and $\mathcal{B} = \{\text{all open intervals } (a, b) : -\infty < a < b < +\infty\}$.

↪ **Proposition 2.4**

Let \mathcal{B} be a collection of open sets in X . Then, \mathcal{B} is a basis \iff

1. $\forall x \in X, \exists U\text{-open} \in \mathcal{B}$ s.t. $x \in U$.
2. If $U_1 \in \mathcal{B}$ and $U_2 \in \mathcal{B}$, and $x \in U_1 \cap U_2$, then $\exists U_3 \in \mathcal{B}$ s.t. $x \in U_3 \subseteq U_1 \cap U_2$.

¹"Requires" proof.

⊗ Example 2.2

Consider $X = \mathbb{R}$. Requirement 1. follows from taking $U = (x - \varepsilon, x + \varepsilon)$ for any $\varepsilon > 0$. For 2., suppose $x \in (a, b) \cap (c, d) =: U_1 \cap U_2$. Let $U_3 = (\max\{a, c\}, \min\{b, d\})$; then, we have that $U_3 \subseteq U_1 \cap U_2$, while clearly $x \in U_3$.

↪ Proposition 2.5

In a metric space, a basis for a topology is a collection of open balls,

$$\{B(x, r) : x \in X, r > 0\} = \{\{y \in X : d(x, y) < r\} : x \in X, r > 0\}.$$

Proof. We prove via proposition 2.4. Property 1. holds clearly; $x \in B(x, \varepsilon)$ -open $\subseteq \mathcal{B}$.

For property 2., let $x \in B(y_1, r_1) \cap B(y_2, r_2)$, that is, $d(x, y_1) < r_1$ and $d(x, y_2) < r_2$. Let

$$\delta := \min\{r_1 - d(x, y_1), r_2 - d(x, y_2)\}.$$

We claim that $B(x, \delta) \subseteq U_1 \cap U_2$.

Let $z \in B(x, \delta)$. Then,

$$d(z, y_1) \stackrel{\Delta\#}{\leq} d(z, x) + d(x, y_1) < \delta + d(x, y_1) \leq r_1 - d(x, y_1) + d(x, y_1) = r_1,$$

hence, as $d(z, y_1) < r_1 \implies z \in B(y_1, r_1) = U_1$. Replacing each occurrence of y_1, r_1 with y_2, r_2 respectively gives identically that $z \in B(y_2, r_2) = U_2$. Hence, we have that $B(x, \delta) \subseteq U_1 \cap U_2$ and 2. holds. ■

2.3 Subspaces

↪ Definition 2.7

Let X be a topological space and let $Y \subseteq X$. We define the subspace topology on Y :

1. Open sets in $Y = \{Y \cap \text{open sets in } X\}$

↪ Proposition 2.6: Consequences of Subspace Topologies

Suppose \mathcal{B} is a basis for a topology in X . Then, $\{U \cap Y : U \in \mathcal{B}\}$ forms a basis for the subspace $Y \subseteq X$.

Suppose X a metric space. Then, Y is also a metric space, with the same distance.

↪ Proposition 2.7

Let $Y \subseteq X$ - a metric space. Then, the metric space topology for (Y, d) is the same as the subspace topology.

Proof. (Sketch) A basis for the open sets in X can be written $\bigcup_{\alpha \in I} B(x_\alpha, r_\alpha)$; hence

$$Y \cap \left(\bigcup_{\alpha \in I} B(x_\alpha, r_\alpha) \right) = \bigcup_{\alpha \in I} (Y \cap B(x_\alpha, r_\alpha))$$

is an open set topology for Y . ■

↪ **Lemma 2.1**

Let $A \subseteq X$ -open, $B \subseteq A$; B -open in subspace topology for $A \iff B$ -open in X .

↪ **Lemma 2.2**

Let $Y \subseteq X, A \subseteq Y$. Then, \bar{A} in $Y = Y \cap \bar{A}$ in X . We can denote this

$$\bar{A}_Y = \bar{A}_X \cap Y.$$

2.4 Continuous Functions

↪ **Definition 2.8: Continuous Function**

Let X, Y be topological spaces. Let $f : X \rightarrow Y$. f is *continuous* $\iff \forall$ open $V \in Y, f^{-1}(V)$ -open in X .

↪ **Proposition 2.8**

This definition is consistent with the normal ε - δ definition on the real line.

Proof. Let $f : \mathbb{R} \rightarrow \mathbb{R}$, continuous; that is, $\forall \varepsilon > 0, \forall x \in \mathbb{R} \exists \delta > 0$ s.t. $|x_1 - x| < \delta$, then $|f(x_1) - f(x)| < \varepsilon$.

Let $V \subseteq \mathbb{R}$ open. Let $y \in V$. Then, $\exists \varepsilon : (y - \varepsilon, y + \varepsilon) \subseteq V$. Let $y = f(x)$, hence $y \in f^{-1}(V)$. Now, if $d(x, x_1) < \delta$, we have that $d(f(x_1), f(x)) < \varepsilon$ (by continuity of f), hence $f(x_1) \in (y - \varepsilon, y + \varepsilon) \subseteq V$; moreover, $(x - \delta, x + \delta) \subseteq f^{-1}(V)$, thus $f^{-1}(V)$ is open as required.

The inverse of this proof follows identically. ■

↪ Lecture 02; Last Updated: Tue Apr 9 21:38:10 EDT 2024

↪ **Proposition 2.9**

Suppose \mathcal{B} forms a basis of topology for Y . Then, $f : X \rightarrow Y$ is continuous if $f^{-1}(U)$ open $\forall U \in \mathcal{B}$.

Proof. If U -open set in Y , then $\exists I$ -index set and a collection of open sets $\{A_\alpha\}_{\alpha \in I}, A_\alpha \in \mathcal{B}$, s.t. $U = \bigcup_{\alpha \in I} A_\alpha$. Then, we have

$$f^{-1}(U) = f^{-1}\left(\bigcup_{\alpha \in I} (A_\alpha)\right) = \bigcup_{\alpha \in I} \underbrace{f^{-1}(A_\alpha)}$$

Hence, if each $f^{-1}(A_\alpha)$ open, then $\cup_{\alpha \in I} f^{-1}(A_\alpha)$ open; hence it suffices to check if $f^{-1}(U) \forall U$ -open in V is open to see if f continuous. ■

↪ **Theorem 2.1: Continuity of Composition**

If $f : X \rightarrow Y$ continuous and $g : Y \rightarrow Z$ continuous, then $g \circ f$ continuous as well.

Proof. Let U -open in Z . Then

$$(g \circ f)^{-1}(U) = \underbrace{f^{-1}(\underbrace{g^{-1}(U)}_{\text{open in } Y})}_{\text{open in } X}$$

↪ **Proposition 2.10**

If $f : X \rightarrow Y$ continuous and $A \subseteq X$, A has subspace topology, then $f|_A : A \rightarrow Y$ is also continuous.²

Proof. Let U -open in Y . Then

$$(f|_A)^{-1}(U) = \underbrace{f^{-1}(U)}_{\text{open}} \cap \underbrace{A}_{\text{open}}$$

By the definition of subspace topology, this is an open set and hence $f|_A$ is continuous. ■

2.5 Product Spaces

↪ **Definition 2.9: Finite Product Spaces**

Let X_1, \dots, X_n be topological spaces. We define

$$(X_1 \times X_2 \times \dots \times X_n),$$

and aim to define a *product topology*; a basis of which consists of cylinder sets.

↪ **Definition 2.10: Cylinder Set**

A *cylinder set* has the form

$$A_1 \times A_2 \times \dots \times A_n$$

where each A_j -open in X_j .

²We denote $f|_A$ as the restriction of the domain of f to A .

⊗ **Example 2.3**

Given an open interval $(a_1, b_1), (a_2, b_2) \subset \mathbb{R}$, the set $(a_1, b_1) \times (a_2, b_2) \subset \mathbb{R}^2$ is a basis for the topology on \mathbb{R}^2 .

↪ **Definition 2.11: Projection**

Let $X_1 \times X_2 \times \cdots \times X_n =: X$. The *projection* $\pi_j : X \rightarrow X_j$ maps $(x_1, \dots, x_n) \rightarrow x_j \in X_j$.

Remark 2.3. One can show π_j continuous.

↪ **Definition 2.12: Coordinate Function**

Given a function $f : Y \rightarrow X_1 \times \cdots \times X_n = (x_1(y), x_2(y), \dots, x_n(y))$. The *coordinate function* is

$$f_j = \pi_j \circ f; \quad f_j = x_j(y).$$

↪ **Proposition 2.11**

$f : Y \rightarrow X = X_1 \times \cdots \times X_n$ continuous $\iff f_j : Y \rightarrow X_j$ continuous.

Proof. Its enough to show that $\forall U \in \mathcal{B}$ -basis for X -product space, $f^{-1}(U)$ -open in Y . Take $U = A_1 \times \cdots \times A_n$ -open. Then, we claim that

$$f^{-1}(U) = f^{-1}(A_1 \times \cdots \times A_n) = f_1^{-1}(A_1) \cap f_2^{-1}(A_2) \cap \cdots \cap f_n^{-1}(A_n). \quad \star$$

If this holds, then as each f_i continuous (being a composition of continuous functions) and each A_i open in X_i , then each $f_i^{-1}(A_i)$ open in Y and hence \star , being the finite intersection of open sets in Y , is itself open in Y . ■

⊗ **Example 2.4: Fourier Transform: Motivation for Infinite Product Topologies**

Let $f \in C([0, 2\pi])$ is real-valued. We write the n th Fourier coefficients

$$\begin{aligned}\hat{f}(n) &= \frac{1}{2\pi} \int_0^{2\pi} f(x)e^{-inx} dx \\ &= \frac{1}{2\pi} \int_0^{2\pi} f(x) \cos(nx) dx - i \frac{1}{2\pi} \int_0^{2\pi} f(x) \sin(nx) dx.\end{aligned}$$

And the Fourier transform of f as the infinite product

$$f(x) \mapsto (\dots, \hat{f}(-n), \hat{f}(-n+1), \dots, \hat{f}(-1), \hat{f}(0), \hat{f}(1), \dots, \hat{f}(n), \dots) \in \prod_{n \in \mathbb{Z}} (\mathbb{C})_n.$$

Hence, this is an (countably, as indexed by integers) infinite product space.

Now, let $f : \mathbb{R} \rightarrow \mathbb{R}$. Suppose $f(x) \rightarrow 0$ “fast enough” as $|x| \rightarrow \infty$ and f continuous. Then, we can define the Fourier coefficients

$$\hat{f}(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} f(x)e^{-itx} dx,$$

where $t \in \mathbb{R}$. We then have the transform

$$f \mapsto \{\hat{f}(t)\}_{t \in \mathbb{R}}.$$

In this case, our index set is \mathbb{R} is (uncountably) infinite.

↪ **Definition 2.13: Product Topology/Cylinder Sets for ∞ Products**

Let $X = \prod_{\alpha \in I} X_\alpha$. Then, a basis for X is given by cylinder sets of the form $A = \prod_{\alpha \in I} A_\alpha$ where A_α -open in X_α , AND $A_\alpha = X_\alpha$ except for finitely many indices α .

That is, there exists a finite set $J = (\alpha_1, \dots, \alpha_k) \subseteq I$, such that we can write $A = \prod_{\alpha \in J} A_\alpha \times \prod_{\alpha \notin J} X_\alpha$ (where A_α open in X_α).

↪ **Proposition 2.12**

Given $f : Y \rightarrow \prod_{\alpha \in I} X_\alpha = X$, then (taking $f_\alpha = \pi_\alpha \circ f$ as before) we have that f is continuous in $X \iff f_\alpha : Y \rightarrow X_\alpha$ continuous in $X_\alpha \forall \alpha \in I$.

Remark 2.4. Extension of proposition 2.11 to infinite product space.

Proof. Write $U = \prod_{\alpha \in J} A_\alpha \times \prod_{\alpha \notin J} X_\alpha$. Then,

$$f^{-1}(U) = \bigcap_{\alpha \in J} f_\alpha^{-1}(A_\alpha)$$

which is open in Y , hence f continuous. ■

Remark 2.5. *The intersection of the entire spaces give no restriction.*

↪ Lecture 03; Last Updated: Fri Jan 19 11:49:27 EST 2024

2.6 Metrizable

↪ Proposition 2.13

Different metrics can define the same topology.

⊗ Example 2.5

1. Different ℓ_p metrics in \mathbb{R}^n (PSET 1)
2. Let (X, d) be a metric space. Then,

$$\tilde{d}(x, y) := \frac{d(x, y)}{d(x, y) + 1}$$

is also a metric (the first two axioms are trivial), and defines the same topology. Note, moreover, that $\tilde{d}(x, y) \leq 1 \forall x, y$; this distance is bounded, and can often be more convenient to work with in particular contexts.

↪ Question 2.1

Suppose (X_k, d_k) are metric spaces $\forall k \geq 1$. Then, we can define the product topology τ on

$$X := \prod_{k=1}^{\infty} X_k.$$

Does the product topology τ come from a metric? That is, is τ metrizable?

Remark 2.6. *There do indeed exist examples of non-metrizable topological spaces; this question is indeed well-founded.*

Answer. Let $\underline{x} = (x_1, x_2, \dots, x_n, \dots)$, $\underline{y} = (y_1, y_2, \dots, y_n, \dots) \in \prod_{k=1}^{\infty} X_k$ (where $x_i, y_i \in X_i$) be infinite sequences of elements. Then, for each metric space X_k take the metric

$$\tilde{d}_k(x_k, y_k) = \frac{d_k(x_k, y_k)}{1 + d_k(x_k, y_k)}$$

(as in the example above). Then, we define

$$D(\underline{x}, \underline{y}) = \sum_{k=1}^{\infty} \frac{\tilde{d}_k(x_k, y_k)}{2^k},$$

noting that $D(\underline{x}, \underline{y}) \leq \sum_{k=1}^{\infty} \frac{1}{2^k} = 1$ (by our construction, “normalizing” each metric), hence this is a valid, *converging* metric (which wouldn’t otherwise be guaranteed if we didn’t normalize the metrics). It remains to show whether this metric omits the same topology as τ . ■

2.7 Compactness, Connectedness

↪ Definition 2.14: Compact

A set A in a topological space is said to be *compact* if every cover has a finite subcover. That is, if

$$A \subseteq \bigcup_{\alpha \in I} U_{\alpha} - \text{open},$$

then $\exists \{\alpha_1, \dots, \alpha_n \in I\}$ such that $A \subseteq \bigcup_{i=1}^n U_{\alpha_i}$.

↪ Proposition 2.14

A closed interval $[a, b]$ is compact.

Proof. If³ $a = b$, this is clear. Suppose $a < b$, and let $[a, b] \subseteq \bigcup_{i \in I} U_i =: \mathcal{U}$ be an arbitrary cover. Then, we proceed in the following steps:

1. **Claim:** Given $x \in [a, b], x \neq b, \exists y \in [a, b]$ s.t. $[x, y]$ has a finite subcover.

Let $x \in [a, b], x \neq b$. Then, $\exists U_{\alpha} \in \mathcal{U} : x \in U_{\alpha}$. Since U_{α} open, and $x \neq b$, we further have that $\exists c \in [a, b]$ s.t. $[x, c] \subseteq U_{\alpha}$.

Now, let $y \in (x, c)$; then, the interval $[x, y] \subseteq [x, c] \subseteq U_{\alpha}$, that is, $[x, y]$ has a finite subcover.

2. Define $C := \{y \in [a, b] : y > a, [a, y] \text{ has a finite subcover}\}$. We note that

- $C \neq \emptyset$; taking $x = a$ in Step 1. above, we have that $\exists y \in [a, b]$ such that $[a, y]$ has a finite step cover, so this $y \in C$.
- C bounded; by construction, $\forall y \in C, a < y \leq c$.

Thus, we can validly define $c := \sup C$, noting that $a < c \leq b$. Ultimately, we wish to prove that $c = b$, completing the proof that $[a, b]$ has a finite subcover.

3. **Claim:** $c \in C$.

Let $U_{\beta} \in \mathcal{U} : c \in U_{\beta}$. Then, by the openness of $U_{\beta}, \exists d \in [a, b]$ s.t. $(d, c] \subseteq U_{\beta}$.

³This proof is adapted from that of Theorem 27.1 in Munkre’s *Topology*, an identical theorem but applied to more general ordered topologies.

Supposing $c \notin C$, then $\exists z \in C$ such that $z \in (d, c)$; if one did not exist, then this would imply that d was a smaller upper bound than c , a contradiction. Thus, $[z, c] \subseteq (d, c) \subseteq U_\beta$.

Moreover, we have that, given $z \in C$, $[a, z]$ has a finite subcover; call it $U_z \subseteq \mathcal{U}$. This gives, then:

$$[a, c] = [a, z] \cup [z, c] \subseteq U_z \cup U_\beta.$$

But this is a finite subcover of $[a, c]$, contradicting the fact that $c \notin C$. We conclude, then, that $c \in C$ after all.

4. **Claim:** $c = b$.

Suppose not; then, since we have $c \leq b$, then assume $c < b$. Then, applying Step 1. with $x = c$ (which we can do, by our assumption of $c \neq b$!), then we have that $\exists y > c$ s.t. $[c, y]$ has a finite subcover, call this $U_y \subseteq \mathcal{U}$.

Moreover, we had $c \in C$, hence $[a, c]$ has a finite subcover, call this $U_c \subseteq \mathcal{U}$.

Then, this gives us that

$$[a, y] = [a, c] \cup [c, y] \subseteq U_c \cup U_y,$$

that is, $[a, y]$ has a finite subcover, and so $y \in C$. But recall that $y > c$; hence, this a contradiction to c being the least upper bound of C . We conclude that $c = b$, and thus $[a, b]$ has a finite subcover, and is thus compact. ■

Remark 2.7. A similar proof shows that $[a, b]$ is connected; we cannot cover it by two disjoint open sets.

↔ **Theorem 2.2: On Compactness**

Let $A \subseteq \mathbb{R}^n$. Then, A compact $\iff A$ closed and bounded.

↔ **Proposition 2.15**

If X, Y are compact topological spaces, then $X \times Y$ is compact.

Remark 2.8. By induction, if X_1, \dots, X_n compact, so is $\prod_{i=1}^n X_i$.

↔ **Proposition 2.16**

A closed subset of a compact topological space is compact in the subspace topology.

Proof. (Of theorem 2.2)

(\Leftarrow) If $A \subseteq \mathbb{R}^n$ closed and bounded, then $A \subseteq [-R, +R]^n$ for some $R > 0$ (it is contained in some “ n -cube”). Then, we have that $[-R, R]$ is compact, by proposition 2.14, proposition 2.15, and proposition 2.16, A itself compact.

(\Rightarrow) Suppose $A \subseteq \mathbb{R}^n$ is compact. Then, $\bigcup_{x \in A} B(x, \varepsilon)$ for some $\varepsilon > 0$ is an open cover of A . As A compact, there must exist a finite subcover of this cover, $A \subseteq \bigcup_{i=1}^N B(x_i, r_i)$. Let $R := \max_{i=1}^N (||x_i|| + r_i)$. Then, $A \subseteq \overline{B(0, R)}$, that is, A is bounded.

Now, suppose x is a limit point of A . Then, any neighborhood of x contains a point in A , so $\forall r > 0, B(x, r) \cap A \neq \emptyset$, and so $\overline{B}(x, r)$ also contains a point of A for any $r > 0$.

Now, suppose $x \notin A$ (looking for a contradiction). Then,

$$U := \bigcup_{r>0} U_r := \bigcup_{r>0} (\mathbb{R}^n \setminus \overline{B}(x, r)) = \mathbb{R}^n \setminus \{x\}$$

is an open cover for the set A . A being compact implies that U has an finite subcover such that $A \subset U_{r_1} \cup U_{r_2} \cup \dots \cup U_{r_N}$. Let $r_0 = \min_{i=1}^N r_i$. Then, $A \subseteq U_{r_0}$, and $A \cap B(x, r_0) = \emptyset$; but this is a contradiction to the definition of a limit point, hence any limit point x is contained in A and A is thus closed by definition. ■

↪ **Proposition 2.17**

Compact \implies sequentially compact; that is, every sequence in a compact set has a convergent subsequence.

↪ Lecture 04; Last Updated: Tue Apr 9 14:45:17 EDT 2024

↪ **Definition 2.15: Connected**

A topological space X is *not connected* if $X = U \cup V$ for two open, nonempty, disjoint sets U, V .

If this does not hold, X is said to be *connected*.

A set $A \subseteq X$ is not connected if A is not connected in the subspace topology $\iff A = \subseteq U \cup V$, for U, V -open in X , $(U \cap A) \neq \emptyset$, $(V \cap A) \neq \emptyset$ and $U \cap V = \emptyset$.

↪ **Theorem 2.3**

Let X be a connected topological space. Let $f : X \rightarrow Y$ be a continuous function. Then, $f(X)$ is also connected.

Proof. Suppose, seeking a contradiction, that X is connected, but $f(X)$ is not. Then, we can write $f(X) \subseteq Y$ as $f(X) \subseteq U \cup V$, such that U, V open in Y and $U \cap V = \emptyset$. Then,

$$(U \cap f(X)) \cap (V \cap f(X)) = \emptyset.$$

We also have that

$$X \subseteq \underbrace{f^{-1}(U)}_{\text{open in } X, \neq \emptyset} \cup \underbrace{f^{-1}(V)}_{\text{open in } X, \neq \emptyset}.$$

$f^{-1}(U) \cap f^{-1}(V) = \emptyset$ (that is, they are disjoint) by our assumption; this is a contradiction to the connectedness of X , as we are able to write it as a subset of two disjoint open sets. Hence, $f(X)$ is indeed connected. ■

↪ **Lemma 2.3**

Any interval $(a, b), [a, b], [a, b), \dots, \subseteq \mathbb{R}$ is connected.

Proof. ■

↪ **Theorem 2.4: “Intermediate Value Theorem”**

Suppose X is connected and $f : X \rightarrow \mathbb{R}$ is a continuous function. Then, f takes intermediate values.

More precisely, let $a = f(x), b = f(y)$ for $x, y \in X$. Assume $a < b$. Then, $\forall a < c < b, \exists z \in X$ s.t. $f(z) = c$.

Proof. Suppose, seeking a contradiction, that $\exists c : a < c < b$ s.t. $c \notin f(X)$ (that is, there exists an intermediate value that is “not reached” by the function).

Let $U = (-\infty, c)$ and $V = (c, +\infty)$; note that these are disjoint open sets. Then, we have that

$$X = f^{-1}(U) \cup f^{-1}(V),$$

by our assumption of $c \notin f(X)$. But this gives that X is not connected, as the union of two open (by continuity), disjoint, nonempty ($f(x) = a \in U \implies x \in f^{-1}(U)$, and $f(y) = b \in V \implies y \in f^{-1}(V)$) sets, a contradiction. ■

↪ **Theorem 2.5**

Suppose X is compact, Y -topological space, $f : X \rightarrow Y$ is a continuous function. Then, $f(X)$ is also compact.

Proof. Let $\{U_\alpha\}_{\alpha \in I}$ be an open cover of $f(X) \subseteq Y$, that is,

$$f(X) \subseteq \bigcup_{\alpha \in I} U_\alpha \implies X \subseteq f^{-1}\left(\bigcup_{\alpha \in I} U_\alpha\right) = \bigcup_{\alpha \in I} f^{-1}(U_\alpha) =: \bigcup_{\alpha \in I} V_\alpha - \text{open.}$$

Then, this is an open cover of X ; X is compact, thus there exists a finite subcover, that is, indices $\{\alpha_1, \dots, \alpha_n\} \subseteq I$ such that $X = \bigcup_{i=1}^n V_{\alpha_i}$. Thus,

$$f(X) \subseteq \bigcup_{i=1}^n U_{\alpha_i},$$

which is a finite subcover of $f(X)$. Thus, $f(X)$ is compact. ■

Remark 2.9. Recall the “extreme value theorem”: let $f : [a, b] \rightarrow \mathbb{R}$ a continuous function; then, a minimum and maximum is obtained for $f(x)$ on this interval for values in this interval.

↪ **Theorem 2.6**

Let X compact, and $f : X \rightarrow \mathbb{R}$ a continuous function. Then,

$$\max_{x \in X} f(x) \text{ and } \min_{x \in X} f(x)$$

are both attained.

Proof. $f(X) \subseteq \mathbb{R}$ is compact by theorem 2.5, and so by theorem 2.2, $f(X)$ is closed and bounded. Let, then, $m = \inf f(X)$ and $M = \sup f(X)$; these necessarily exist, since $f(X)$ is bounded. Both m and M are limit points of $f(X)$. But $f(X)$ is closed, and hence contains all of its limit points, and thus $m \in f(X)$ and $M \in f(X)$, and thus $\exists y_m : f(y_m) = m$ and $y_M : f(y_M) = M$. ■

↪ **Definition 2.16: Path Connected**

A set $A \subseteq X$ is called *path connected* if $\forall x, y \in A, \exists f : [a, b] \rightarrow X$, continuous, s.t. $f(a) = x, f(b) = y$ and $f([a, b]) \subseteq A$.

The set $\{f(t) : a \leq t \leq b\}$ is called a *path* from x to y .

↪ **Theorem 2.7: Path connected \implies connected**

If $A \subseteq X$ is path connected, then A is connected.

Proof. Suppose, seeking a contradiction, that A is path connected, but not connected. Then, we can write $A \subseteq U \cup V$, for open, disjoint, nonempty subsets $U, V \subseteq X$.

Let $x \in U \cap A$ and $y \in V \cap A$. Then, $\exists f : [a, b] \rightarrow A$ s.t. $f(a) = x, f(b) = y$, and $f([a, b]) \subseteq A$, by the path connectedness of A . Then,

$$[a, b] \subseteq f^{-1}(A) \subseteq \underbrace{f^{-1}(U \cap A)}_{\text{open}} \cup \underbrace{f^{-1}(V \cap A)}_{\text{open}} =: \underbrace{U_1}_{a \in} \cup \underbrace{U_2}_{b \in},$$

that is, $[a, b]$ is contained in a union of open, nonempty, disjoint sets, contradicting $[a, b]$ the connectedness of $[a, b]$ by lemma 2.3. Thus, A is connected. ■

Remark 2.10. A counterexample to the opposite side of the implication is the Topologist's sine curve, the set

$$\left\{ \left(x, \sin \left(\frac{1}{x} \right) \right) : x \in (0, 1] \right\} \cup \{0\} \times [-1, 1].$$

This set is connected in \mathbb{R}^2 , but is not path connected.

↪ **Proposition 2.18**

For open sets in \mathbb{R}^n , path connected \iff connected.

2.8 Path Components, Connected Components

Remark 2.11. Remark that if a metric space X is not connected, then we can write $X = U \cup V$ where U, V are open, nonempty and disjoint. It follows, then, that $U = V^c$ (and vice versa) and hence U, V are both open and closed.

↪ Definition 2.17: Connected Component

A connected component of $x \in X$ is the largest connected subset of X that contains x .

⊗ Example 2.6

Let $X = (0, 1) \cup (1, 2)$. Here, we have two connected components, $(0, 1)$ and $(1, 2)$

⊗ Example 2.7: Middle Thirds Cantor Set

Let $C_0 := [0, 1]$, and given C_n , define $C_{n+1} := \frac{1}{3}(C_n \cup (2 + C_n))$ for $n \geq 0$. C_∞ is totally disconnected.

↪ Definition 2.18: Path Component

A path component $P(x)$ of $x \in X$ is the largest path connected subset of X that contains x .

↪ Proposition 2.19

$P(x) = \{x \in X : \exists \text{ continuous path } \gamma : [0, 1] \rightarrow X : \gamma(0) = x, \gamma(1) = y\}$.

Remark 2.12. Where we “start” a path does not matter. We write $x \sim y$ if $\exists \gamma$ from x to y ; this is an equivalence relation on the elements of X .

Remark 2.13. The choice of $[0, 1]$ here is arbitrary; any closed interval is homeomorphic.

↪ Lemma 2.4

If $P(x) \cap P(y) \neq \emptyset$, then $P(x) = P(y)$.

Proof. $P(x) \cap P(y) \neq \emptyset \implies \exists z : x \sim z \wedge y \sim z \implies x \sim y$. ■

↪ Lemma 2.5

If $A \subseteq X$ is connected, then \bar{A} is also connected.

↪ Lemma 2.6

Suppose $A \subseteq X$ is both open and closed. Then, if $C \subseteq X$ is connected and $C \cap A \neq \emptyset$, then $C \subseteq A$.

Proof. If A is both open and closed, then $C \cap A$ is both open and closed in C . If $C \cap A^C \neq \emptyset$, then this is also open and closed in C . Hence, we can write $C = (C \cap A) \cup (C \cap A^C)$, that is, a disjoint union of two nonempty open sets, contradicting the connectedness of C . Hence, $C \cap A^C = \emptyset$, and so $C \subseteq A$. ■

↪ **Proposition 2.20**

Let $\{C_\alpha\}_{\alpha \in I}$ be a collection of nonempty connected subspaces of X s.t. $\forall \alpha, \beta \in I, C_\alpha \cap C_\beta \neq \emptyset$. Then, $\bigcup_{\alpha \in I} C_\alpha$ is connected.

↪ **Proposition 2.21**

Suppose each $x \in X$ has a path-connected neighborhood. Then, the path components in X are the same as the connected components in X .

2.8.1 Cantor Staircase Function

↪ **Definition 2.19: An Explicit Definition**

Let $x \in C : x = 0.a_1a_2a_3 \dots$ (base 3), ie $a_j = \begin{cases} 0 \\ 2 \end{cases}$. Define

$$f(x) = \begin{cases} \sum \frac{a_j/2}{2^j} & x \in C \\ \text{extend by continuity} & x \notin C. \end{cases}$$

That is, if $x \notin C$, set $f(y) = \sup_{x \in C, x < y} f(x) = \inf_{x \in C, x > y} f(x)$.

↪ **Definition 2.20: Complement Definition**

To construct the complement of the Cantor set, begin with $[0, 1]$ and at a step n , we remove 2^n open intervals from this interval. $f(x)$ will be constant on each of these intervals with values $\frac{k}{2^n}$ where k odd and $0 < k < 2^n$. Extend by continuity to all $x \in C$.

Remark 2.14. *Wikipedia's explanation of this is far better than whatever this definition is trying to say.*

3 L^p SPACES

3.1 Review of ℓ^p Norms

Remark 3.1. Recall that for $1 \leq p \leq +\infty$, we define for $x = (x_1, \dots, x_n) \in \mathbb{R}^n$ the norm

$$\|x\|_p = (|x_1|^p + \dots + |x_n|^p)^{\frac{1}{p}}, \quad \|x\|_\infty = \max_{i=1}^n |x_i|.$$

Similarly, for infinite vector spaces, we had, for $x = (x_1, \dots, x_n, \dots)$, the norm

$$\|x\|_p = \left(\sum_{i=1}^{\infty} |x_i|^p \right)^{\frac{1}{p}}, \quad \|x\|_\infty = \sup_{i \geq 1} |x_i|.$$

Here, we define

$$\ell_p := \{x = (x_1, \dots, x_n) : \|x\|_p < +\infty\}.$$

3.2 ℓ^p Norms, Hölder-Minkowski Inequalities

↪ Definition 3.1: Hölder Conjugates

For $1 \leq p, q \leq +\infty$, we say that p, q are said to be Hölder conjugates if

$$\frac{1}{p} + \frac{1}{q} = 1.$$

Remark 3.2. We refer to these simply as “conjugates” throughout as no other conception of conjugate numbers will be discussed.

Further, we take by convention $\frac{1}{\infty} = 0$.

↪ Proposition 3.1: Hölder’s Inequality

Let $x = (x_1, \dots, x_n), y = (y_1, \dots, y_n) \in \mathbb{R}^n$. Suppose $p, q : 1 \leq p, q \leq +\infty$ are conjugate. Then,

$$\langle x, y \rangle_{\mathbb{R}^n} := \left| \sum_{i=1}^n x_i y_i \right| \leq \|x\|_p \cdot \|y\|_q$$

⊗ Example 3.1

For the case $p = 1$ or ∞ (functionally, the same case):

↪ **Lemma 3.1**

Let p, q be conjugates, and $x, y \geq 0$. Then,

$$xy \leq \frac{x^p}{p} + \frac{y^q}{q}.$$

Remark 3.3. If the inequality holds, then, for some $t > 0$, let $\tilde{x} = t^{\frac{1}{p}} \cdot x$, $\tilde{y} = t^{\frac{1}{q}} y$. Substituting x for \tilde{x} and y for \tilde{y} , we have

$$\text{LHS: } \tilde{x}\tilde{y} = t^{\frac{1}{p}}x \cdot t^{\frac{1}{q}}y = t^{\frac{1}{p}+\frac{1}{q}} \cdot xy = xy$$

$$\text{RHS: } \dots = t\left(\frac{x^p}{p} + \frac{y^q}{q}\right).$$

That is, we have

$$t \cdot xy \leq t \left(\frac{x^p}{p} + \frac{y^q}{q} \right),$$

hence, the inequality is preserved under multiplication by a positive scalar; moreover, the original inequality holds iff this “scaled” version holds. Hence, choosing t such that $\tilde{y} = 1$ (let $t = \left(\frac{1}{y}\right)^q$), it suffices to prove the lemma for $y = 1$.

Proof. If $x = 0$ or $y = 0$, then the entire LHS becomes 0 and we are done; assume $x, y > 0$; by the previous remark, assume wlog $y = 1$. Then, we have

$$\begin{aligned} x \cdot y \leq \frac{x^p}{p} + \frac{y^q}{q} &\iff x \cdot 1 \leq \frac{x^p}{p} + \frac{1}{q} \\ &\iff \frac{x^p}{p} - x + \frac{1}{q} =: f(x) \geq 0. \end{aligned}$$

Taking the derivative, we have

$$\begin{aligned} f'(x) &= \frac{p x^{p-1}}{p} - 1 = x^{p-1} - 1 \\ p > 1 &\implies p - 1 > 0 \implies \begin{cases} f'(x) > 0 & \forall x > 1 \\ f'(x) = 0 & x = 0 \\ f'(x) < 0 & \forall 0 < x < 1 \end{cases} \end{aligned}$$

Hence, $x = 1$ is a local minimum of the function, and thus $f(x) \geq f(1) \forall 0 < x \leq 1$. But $f(1) = \frac{1^p}{p} - 1 + \frac{1}{q} = 1 - 1 = 0$, hence $f(x) \geq 0 \forall x \geq 0$, as desired, and the inequality holds. ■

Proof. Assume $\|x\|_p = \|y\|_q = 1$. Then,

$$\begin{aligned}
 \left| \sum_{i=1}^n x_i y_i \right| &\leq \sum_{i=1}^n |x_i y_i| && \text{(by triangle inequality)} \\
 &\leq \sum_{i=1}^n \left| \frac{x_i^p}{p} + \frac{y_i^q}{q} \right| && \text{(by lemma 3.1)} \\
 &= \frac{1}{p} \left(\sum_{i=1}^n |x_i|^p \right) + \frac{1}{q} \left(\sum_{i=1}^n |y_i|^q \right) \\
 &= \frac{1}{p} \|x\|_p^p + \frac{1}{q} \|y\|_q^q && \text{(by starring)} \\
 &= \frac{1}{p} \cdot 1^p + \frac{1}{q} \cdot 1^q = \frac{1}{p} + \frac{1}{q} = 1 && \text{(by assumption)} \\
 &= \|x\|_p \cdot \|y\|_q,
 \end{aligned}$$

and the proposition holds, in the special case $\|x\|_p = \|y\|_q = 1$.

If $\|x\|_p = 0$ or $\|y\|_q = 0$, then $x_1 = \dots = x_n = 0$ or $y_1 = \dots = y_n = 0$, resp, then we'd have ($\|x\|_p = 0$ case)

$$0 \cdot y_1 + \dots + 0 \cdot y_n \leq 0,$$

which clearly holds.

Assume, then, $\|x\|_p > 0, \|y\|_q > 0$. Let $\tilde{x} := \frac{x}{\|x\|_p}, \tilde{y} := \frac{y}{\|y\|_q}$. Then,

$$\|\tilde{x}\|_p^p = \frac{(\sum_{i=1}^n |x_i|^p)}{\|x\|_p^p} = \frac{\|x\|_p^p}{\|x\|_p^p} = 1 \implies \|\tilde{x}\|_p = 1.$$

The same case holds for \tilde{y} , hence $\|\tilde{y}\|_q = 1$; that is, we have "rescaled" both vectors. Hence, we can use the case we proved above for when the norms were identically 1 on \tilde{x}, \tilde{y} . We have:

$$\left| \sum_{i=1}^n \tilde{x}_i \tilde{y}_i \right| \leq 1$$

But by definition of \tilde{x}, \tilde{y} , we have

$$\left| \sum_{i=1}^n \tilde{x}_i \tilde{y}_i \right| = \left| \frac{1}{\|x\|_p \|y\|_q} \sum_{i=1}^n x_i y_i \right| \leq 1 \implies \left| \sum_{i=1}^n x_i y_i \right| \leq \|x\|_p \cdot \|y\|_q,$$

and the proof is complete. ■

↪ **Proposition 3.2: Minkowski Inequality**

Let $1 \leq p \leq \infty$, $x, y \in \mathbb{R}^n$. Then,

$$\|x + y\|_p \leq \|x\|_p + \|y\|_p.$$

Remark 3.4. This is just the triangle inequality for ℓ_p norms.

Proof. The cases $p = 1, \infty$ are left as an exercise.

Assume $1 < p < \infty$. Then,

$$\begin{aligned} \|x + y\|_p^p &= \sum_{j=1}^n |x_j + y_j|^p = \sum_{j=1}^n |x_j + y_j| |x_j + y_j|^{p-1} \\ &\leq \sum_{j=1}^{\infty} (|x_j| + |y_j|) \cdot |x_j + y_j|^{p-1} \\ &= \underbrace{\sum_{j=1}^n |x_j| \cdot |x_j + y_j|^{p-1}}_{:=A} + \overbrace{\sum_{j=1}^n |y_j| \cdot |x_j + y_j|^{p-1}}{:=B} \quad \circledast \end{aligned}$$

Let $\vec{u} = (|x_1|, \dots, |x_n|)$ and $\vec{v} = (|x_1 + y_1|^{p-1}, \dots, |x_n + y_n|^{p-1})$, then, $A = \vec{u} \cdot \vec{v} = \langle \vec{u}, \vec{v} \rangle_{\mathbb{R}^n}$. We have

$$\begin{aligned} \|\vec{u}\|_p &= \left(\sum_{i=1}^n (|x_i|^p) \right)^{\frac{1}{p}} = \|x\|_p \\ \|\vec{v}\|_q &= \left(\sum_{i=1}^n (|x_i + y_i|^{p-1})^q \right)^{\frac{1}{q}} \\ &= \left[\sum_{i=1}^n (|x_i + y_i|^{p-1})^{\frac{p}{p-1}} \right]^{\frac{p-1}{p}} \\ &= \left[\sum_{i=1}^n |x_i + y_i|^p \right]^{\frac{p-1}{p}} \\ &= \|x + y\|_p^{p-1} \end{aligned}$$

where the second-to-last line follows from p, q being conjugate, hence $q = \frac{p}{p-1}$. Thus, by [Hölder's Inequality](#), we have that

$$A = \langle \vec{u}, \vec{v} \rangle \leq \|\vec{u}\|_p \cdot \|\vec{v}\|_q = \|x\|_p \cdot \|x + y\|_p^{p-1}.$$

By a similar construction, we can show that

$$B \leq \|y\|_p \cdot \|x + y\|_p^{p-1}.$$

Thus, returning to our original inequality in \otimes , we have

$$\begin{aligned} \|x + y\|_p^p &\leq A + B \\ &\leq \|x\|_p \cdot \|x + y\|_p^{p-1} + \|y\|_p \cdot \|x + y\|_p^{p-1} \\ &\implies \|x + y\|_p \leq \|x\|_p + \|y\|_p, \end{aligned}$$

and the proof is complete. ■

↪ Lecture 07; Last Updated: Tue Jan 30 12:54:59 EST 2024

3.3 Complete Metric Spaces, Completeness of ℓ_p

↪ Theorem 3.1

The sequence of centers of balls with monotonically decreasing radii is a Cauchy sequence in X .

Proof. Let $\varepsilon > 0$ and let $N : \forall j > N, r_j < \varepsilon$. Then,

$$d(x_j, x_k) < r_{\min(j,k)} = r_j$$

↪ Definition 3.2: Complete Metric Space

A metric space is complete if every Cauchy sequence converges to a limit in that space.

\otimes Example 3.2: Examples of Complete Metric Spaces

1. \mathbb{R} , p -adic integers (\mathbb{Z}_p)/rationals (\mathbb{Q}_p).
2. $\ell_p = \{x = (x_i)_{i=1}^{\infty} : \sum_{i=1}^{\infty} |x_i|^p < +\infty\}$, $1 \leq p < +\infty$
3. $\ell_{\infty} = \{x = (x_i) : \sup_{i=1}^{\infty} |x_i| < +\infty\}$.

↪ Proposition 3.3

Hölder's Inequality and Minkowski Inequality inequalities hold for infinite sequences. that is,

1. if $x = (x_i) \in \ell_p$ and $y = (y_i) \in \ell_q$ with $\frac{1}{p} + \frac{1}{q} = 1$, then

$$\left| \sum_{i=1}^{\infty} x_i y_i \right| \leq \|x\|_{\ell_p} \|y\|_{\ell_q}.$$

2. if $x, y \in \ell_p$, then

$$\|x + y\|_p \leq \|x\|_p + \|y\|_p.$$

Remark 3.5. 2. gives the triangle inequality for the $\|x\|_p$ norm on ℓ_p .

Moreover,

$$\begin{aligned} \|c \cdot x\|_p &= \|(c_1x_1, \dots, c_nx_n, \dots)\|_p \\ &= \left(\sum_{i=1}^{\infty} |cx_i|^p \right)^{\frac{1}{p}} = \left(\sum_{i=1}^{\infty} |c|^p |x_i|^p \right)^{\frac{1}{p}} \\ &= (|c|^p)^{\frac{1}{p}} \|x\|_p = c \cdot \|x\|_p \end{aligned}$$

Proof. (of 2.) If $x, y \in \ell_p$, we have that $\sum_{i=1}^{\infty} |x_i|^p < +\infty, \sum_{i=1}^{\infty} |y_i|^p < +\infty$, so $\exists N > 0 : \sum_{i=N+1}^{\infty} |x_i|^p < \varepsilon, \sum_{i=N+1}^{\infty} |y_i|^p < \varepsilon$. Let $x_i^{(n)} = (x_1, \dots, x_n, 0, 0, \dots)$ be (x) truncated after n (finite) coordinates. This gives

$$\|(x_i + y_i)^{(n)}\|_p \leq \|x_i^{(n)}\|_p + \|y_i^{(n)}\|_p \leq \|x\|_p + \|y\|_p$$

by Minkowski on finite spaces. Taking $n \rightarrow \infty$ (ie, “detruncating”), we have $(x + y) \in \ell_p$, and thus $\|x + y\|_p \leq \|x\|_p + \|y\|_p$.

1. left as an exercise. ■

↔ Proposition 3.4

Let $1 \leq p \leq +\infty$, and $\|x\|_{\infty} = \sup_{i=1}^{\infty} |x_i| = A < +\infty, \|y\|_{\infty} = \sup_{i=1}^{\infty} |y_i| = B < +\infty$. Then, the triangle inequality $\|x + y\|_{\infty} \leq \|x\|_{\infty} + \|y\|_{\infty}$ holds.

Proof. We have

$$\sup_{i=1}^{\infty} |x_i + y_i| \leq \sup_{i=1}^{\infty} (|x_i| + |y_i|) \leq \sup_{i=1}^{\infty} |x_i| + \sup_{i=1}^{\infty} |y_i| = \|x\|_{\infty} + \|y\|_{\infty}.$$

↔ Proposition 3.5

$\|x\|_{\infty} := \sup_{i=1}^{\infty} |x_i|$ is a well-defined norm on ℓ_{∞} .

Proof. The triangle inequality is prove in proposition 3.4. The remainder of the requirements are left as an exercise. ■

↔ Proposition 3.6

$\ell_p \subseteq \ell_q$ if $p < q$.

Proof. Let $x \in \ell_p$. If $\sum_{i=1}^{\infty} |x_i|^p < +\infty$, then $\exists N : \forall i \geq N, |x_i| \leq 1$. Then,

$$\begin{aligned} \sum_{i \geq N} |x_i|^q &\leq \sum_{i \geq N} |x_i|^p < \infty \\ \implies \sum_{i=1}^{\infty} |x_i|^q < +\infty &\implies x \in \ell_q \\ &\implies \ell_p \subseteq \ell_q \end{aligned}$$

■

↪ Lecture 08; Last Updated: Thu Mar 28 09:13:10 EDT 2024

3.4 Contraction Mapping Theorem

↪ Definition 3.3: Contraction Mapping

Let (X, d) be a metric space. A *contraction mapping* on X is a function $f : X \rightarrow X$ for which \exists a constant $0 < c < 1$ such that

$$d(f(x), f(y)) \leq c \cdot d(x, y) \quad \forall x, y \in X.$$

↪ Theorem 3.2: Contraction Mapping Theorem

Let (X, d) be a complete metric space, and let $f : X \rightarrow X$ be a contraction. Then, there exists a unique fixed point z of f such that $f(z) = z$.

Moreover, $f^{[n]}(x) := f \circ f \circ \dots \circ f(x) \rightarrow z$ as $n \rightarrow \infty$ for any $x \in X$.

Remark 3.6. The “functional construction” of the Cantor set is an example of a contraction mapping, with $f_1(x) = \frac{x}{3}$, $f_2(x) = \frac{x+2}{3}$. The first has a fixed point of 0, and the second a fixed point of 1.

Remark 3.7. This is a generalization of [this proof](#) done in Analysis I, an equivalent claim over the reals.

Proof. Fix $x \in X$. Consider the sequence $\{x_0, x_1, x_2, \dots, x_n, \dots\} := \{x, f(x), f \circ f(x), \dots, f^{[n]}(x), \dots\}$ (we call $f^{[n]}$ the *orbit* of x under iterations of f). We claim that this is a Cauchy sequence. Let $n \in \mathbb{N}$ arbitrary, then we have, by the property of the contraction mapping,

$$d(f^{[n+1]}(x) - f^{[n]}(x)) \leq c \cdot d(f^{[n]}(x) - f^{[n-1]}(x)) \leq c^2 d(f^{[n-1]}(x) - f^{[n-2]}(x)).$$

Arguing inductively, it follows that

$$d(f^{[n+1]}(x) - f^{[n]}(x)) \leq c^n d(f(x), x). \quad \star$$

Let now $m, k \in \mathbb{N}, m, k > 0$. It follows that

$$\begin{aligned} d(f^{[m]}, f^{[m+k]})(x) &\leq d(f^{[m]}(x), f^{[m+1]}(x)) + d(f^{[m+1]}(x), f^{[m]}(x)) + \dots + d(f^{[m+k-1]}(x), f^{[m+k]}(x)) \\ &\stackrel{\star}{\leq} d(x, f(x)) [c^m + c^{m+1} + \dots + c^{m+k-1}] \\ &\leq c^m d(x, f(x)) [1 + c + \dots + c^k + c^{k+1} + \dots] = \frac{c^m d(x, f(x))}{1 - c} \end{aligned}$$

Now, given $\varepsilon > 0$, choose N such that $\frac{c^N d(x, f(x))}{1 - c} < \varepsilon$. It follows, then, that $\{f^{[n]}(x)\}_{n \in \mathbb{N}}$ a Cauchy sequence, and thus converges, $f^{[n]}(x) \rightarrow z$ as $n \rightarrow \infty$ for some z .

We further have to show that $f(z) = z$. It is easy to show that f continuous due to the contraction mapping (it is clearly Lipschitz with constant c), and it thus follows that

$$\lim_{n \rightarrow \infty} f(f^{[n]}(x)) = \lim_{n \rightarrow \infty} f^{[n]}(x) \implies f(z) = z,$$

by sequential characterization of continuous functions.

Finally, we need to show that this limit is unique. Suppose $\exists y_1 \neq y_2$, ie two fixed points with $f(y_1) = y_1$ and $f(y_2) = y_2$. Then, by the property of the contraction mapping,

$$d(f(y_1), f(y_2)) \leq c \cdot d(y_1, y_2),$$

but by assumption of being fixed points,

$$d(f(y_1), f(y_2)) = d(y_1, y_2),$$

implying $d(y_1, y_2) \leq c \cdot d(y_1, y_2)$. This is only possible if $d(y_1, y_2) = 0$, and thus $y_1 = y_2$ and the fixed point is indeed unique. ■

↪ Theorem 3.3: ℓ_p complete

The space ℓ_p is complete for all $1 \leq p \leq +\infty$.

Equivalently, if $(x^1), (x^2), \dots, (x^n)$ is a Cauchy sequence in ℓ^p , $\exists y \in \ell^p$ s.t. $x^n \rightarrow y$ as $n \rightarrow \infty$.

Proof. (Sketch) We suppose first $p < +\infty$. Consider an arbitrary number of Cauchy sequences in ℓ_p :

$$\begin{aligned} x^{(1)} &= (x_1^{(1)}, \dots, x_n^{(1)}, \dots) \\ x^{(2)} &= (x_1^{(2)}, \dots, x_n^{(2)}, \dots) \\ &\vdots \quad \quad \quad \vdots \\ x^{(k)} &= (x_1^{(k)}, \dots, x_n^{(k)}, \dots) \in \ell_p \end{aligned}$$

We claim that, for any $k \in \mathbb{N}$, the $(x_k^{(n)})_{n \in \mathbb{N}}$ is a Cauchy sequence; note that in this definition we are taking a *fixed-index* (namely, the k th) element from different sequences (namely, the n th sequence).

Since $x^{(1)}, x^{(2)}, \dots, x^{(n)}, \dots$ are Cauchy sequences in ℓ^p , we have for a fixed $\varepsilon > 0$, $\exists N \in \mathbb{N} : \forall m, n > N$, $d_p(x^{(m)}, x^{(n)}) < \varepsilon$:

$$\begin{aligned} d_p(x^{(m)}, x^{(n)})^p &= \|x^{(m)} - x^{(n)}\|_p^p = \sum_{i=1}^{\infty} |x_i^{(m)} - x_i^{(n)}|^p < \varepsilon^p \\ |x_k^{(m)} - x_k^{(n)}|^p &\leq \sum_{i=1}^{\infty} |x_i^{(m)} - x_i^{(n)}|^p \implies |x_k^{(m)} - x_k^{(n)}|^p < \varepsilon^p \\ &\implies |x_k^{(m)} - x_k^{(n)}| < \varepsilon, \end{aligned}$$

since we are taking “less of the summands in the second line”. It follows, then, that for each k , $\exists z_k : x_k^{(n)} \rightarrow z_k$ as $n \rightarrow \infty$. Let $z = (z_1, \dots, z_n, \dots)$. We claim that $x^{(n)} \rightarrow z \in \ell_p$ as $n \rightarrow \infty$.

First, we show that $d_p(x^{(n)}, z) \rightarrow 0$ as $n \rightarrow \infty$ (that is, $x^{(n)} \rightarrow z$ as $n \rightarrow \infty$). Fix $\varepsilon > 0$, and choose $N \in \mathbb{N}$ for which $d_p(x^{(m)}, x^{(n)}) < \varepsilon \forall m, n \geq N$ (by Cauchy). Fix $K \in \mathbb{N}, K > 0$.

$$\begin{aligned} d_p^p(x^{(n)}, z) &= \|x^{(n)} - z\|_p^p = \sum_{i=1}^{\infty} |x_i^{(n)} - z_i|^p \\ \|x^{(m)} - x^{(n)}\|_p^p < \varepsilon^p &\implies \sum_{i=1}^K |x_i^{(m)} - x_i^{(n)}|^p \leq \varepsilon^p \end{aligned}$$

Let $m \rightarrow \infty$; then $x_i^{(m)} \rightarrow z_i$ (note that i fixed!), and we have

$$\sum_{i=1}^K |z_i - x_i^{(n)}|^p \leq \varepsilon^p.$$

Let $K \rightarrow \infty$; then,

$$\sum_{i=1}^{\infty} |z_i - x_i^{(n)}|^p \leq \varepsilon^p \implies \|z - x^{(n)}\|_p \leq \varepsilon \implies d_p(z, x^{(n)}) \leq \varepsilon,$$

and thus $x^{(n)} \rightarrow z$ as $n \rightarrow \infty$.

It remains to show that $z \in \ell_p$, ie $\|z\|_p < +\infty$. We have:

$$\|z\|_p \leq \underbrace{\|z - x^{(n)}\|_p}_{\rightarrow 0} + \|x^{(n)}\|_p.$$

For sufficiently large n , $\|z - x^{(n)}\|_p \leq 1$ (for instance); $x^{(n)} \in \ell_p$, hence $\|x^{(n)}\|_p < +\infty$ (say, $\|x^{(n)}\|_p \leq M$). Thus:

$$\|z\|_p \leq 1 + M < +\infty \implies z \in \ell_p,$$

and the proof is complete. ■

3.5 Equivalent Notions of Compactness in Metric Spaces

↪ Definition 3.4: Totally Bounded

Let (X, d) be a metric space. If for every $\varepsilon > 0$, $\exists x_1, \dots, x_n \in X, n = n(\varepsilon) : \bigcup_{i=1}^n B(x_i, \varepsilon) = X$, we say X is *totally bounded*.

↪ Lecture 09; Last Updated: Tue Apr 9 22:27:24 EDT 2024

↪ Theorem 3.4

Let (X, d) be a metric space. TFAE:

1. X is complete and totally bounded;
2. X is compact;
3. X is sequentially compact (every sequence has a convergent subsequence).

Proof. (1. \implies 2.) Suppose X complete and totally bounded. Assume towards a contradiction that X not compact, ie there exists an open cover $\{U_\alpha\}_{\alpha \in I}$ of X with no finite subcover.

X being totally bounded gives that it can be covered by finitely many open balls of radius $\frac{1}{2}$. It must be that at least one of these open balls cannot be finitely covered, otherwise we would have a finite subcover. Let F_1 be the closure of this ball. F_1 closed, with diameter $\text{diam}(F_1) \leq 1$. X .

We also have that X can be covered by finitely many balls of radius $\frac{1}{4}$; again, there must be at least one ball B_1 such that $B_1 \cap F_1$ cannot be covered by finitely many open sets from the cover. Let $F_2 = \overline{B_1} \cap F_1$ -closed, with $\text{diam}(F_2) \leq \frac{1}{4} + \frac{1}{4} = \frac{1}{2}$.⁴

Arguing inductively, at some step n , X can be covered by finitely many balls of radius $\frac{1}{2^n}$; at least one of these balls B cannot be covered by a finite subcover hence $B \cap F_{n-1}$ cannot be covered by finitely many U_α 's. Let $F_n = \overline{B} \cap F_{n-1}$ -closed, with $\text{diam}(F_n) \leq \frac{1}{2^{n-1}}$.

As such, we have a nested sequence $F_1 \supseteq F_2 \supseteq \dots \supseteq F_n \supseteq \dots$ of closed sets, where $\text{diam}(F_k) \leq \frac{1}{2^{k-1}} \rightarrow 0$ as $k \rightarrow \infty$.

↪ Lemma 3.1 (Cantor Intersection Theorem). $\bigcap_{k=1}^{\infty} F_k \neq \emptyset$.

Proof. (Of Lemma) Let $x_k \in F_k$. Then, $\{x_k\}_{k \in \mathbb{N}}$ is a Cauchy sequence, since

$$d(x_n, x_{n+k}) \leq \text{diam}(F_n) + \dots + \text{diam}(F_{n+k}) \leq \frac{1}{2^{n-1}},$$

by the nested property, which can be made arbitrarily small for sufficiently large n, k . Hence, $x_n \rightarrow y \in X$ for some y , as X complete. The tail of x_n lies in F_n for all sufficiently large n , and as each F_n closed, the limit must lie in F_n for all sufficiently large n . We conclude the intersection nonempty. ■

⁴ B_1 has radius $\frac{1}{4}$ and hence diameter $\frac{1}{2}$. The intersection of B_1 with a set with a larger diameter must have diameter $\leq \frac{1}{2}$.

This y from the lemma is covered by some U_{α_0} -open for some $\alpha_0 \in I$. Being open, $\exists \varepsilon > 0 : B(y, \varepsilon) \subseteq U_{\alpha_0}$. Let $n : \frac{1}{2^{n-1}} < \varepsilon$. Then, $y \in F_n$, and as $\text{diam}(F_n) \leq \frac{1}{2^{n-1}}$, we have that $F_n \subseteq B(y, \frac{1}{2^{n-1}}) \subseteq B(y, \varepsilon) \subseteq U_{\alpha_0}$. But then, we have that F_n covered by a single open set U_{α_0} , a contradiction to our inductive construction of F_n . We conclude X compact.

(2. \implies 3.) Suppose X compact. Let $\{x_n\}_{n \in \mathbb{N}} \in X$. Let $F_n = \overline{\bigcup_{k \geq n} \{x_k\}}$ -closed; we have too that $F_1 \supseteq F_2 \supseteq \dots \supseteq F_n \supseteq \dots$.

\hookrightarrow **Definition 3.5: Finite Intersection Property**

\mathcal{F} has finite intersection property provided any finite subcollection of sets in \mathcal{F} has a non-empty intersection.

\hookrightarrow **Lemma 3.2** (Finite Intersection Formulation of Compactness). X -compact \iff every collection \mathcal{F} of closed subsets of X with finite intersection property has non-empty intersection.

Proof. ■

This lemma directly gives that $\bigcap_{n=1}^{\infty} F_n \neq \emptyset$, $\{F_n\}_{n \in \mathbb{N}}$ being a collection of closed subsets with any subset having nonempty intersection (by the nestedness). Let $y \in \bigcap_{n=1}^{\infty} F_n$. Take $B(y, \frac{1}{k})$, which thus has nonempty intersection with $\{x_k\}_{k \geq n} \forall n$, ie $\exists n_1 : d(y, x_{n_1}) < 1$ and $\exists n_2 > n_1 : d(y, x_{n_2}) < \frac{1}{2}$. Arguing inductively, $\exists n_j > n_{j-1} : d(y, x_{n_j}) < \frac{1}{j}$ for any given n_{j-1} . It follows that $\lim_{j \rightarrow \infty} x_{n_j} = y$, and thus $\{x_{n_j}\}$ is a convergent subsequence of $\{x_n\}$ that converges within X , and thus X is sequentially compact.

(3. \implies 1.) Suppose X sequentially compact. Let $\{x_n\} \in X$ be a Cauchy sequence in X , which thus have a convergent subsequence $\{x_{n_k}\} \rightarrow y$.

\hookrightarrow **Lemma 3.3.** Let $\{x_n\}$ be a Cauchy sequence in X where X sequentially compact. Then, if $\{x_{n_k}\} \rightarrow y$, so does $\{x_n\} \rightarrow y$

Proof. ■

Then, $\{x_n\}_n \rightarrow y$ and so X complete.

Suppose X not totally bounded, ie $\exists \varepsilon > 0 : X$ cannot be covered by a finite union of balls of $B(x_j, \varepsilon)$. Let $x_1 \in X$ s.t. $B(x_1, \varepsilon) \not\supseteq X$; $\exists x_2 \in X \setminus B(x_1, \varepsilon)$, and so $X \not\subseteq B(x_1, \varepsilon) \cup B(x_2, \varepsilon)$ by assumption. Then, choose $x_3 \in X \setminus (B(x_1, \varepsilon) \cup B(x_2, \varepsilon))$. Arguing inductively, we have that $\exists x_n \in X \setminus (\bigcup_{i=1}^n B(x_i, \varepsilon))$, noting that $d(x_n, x_j) \geq \varepsilon \forall 1 \leq j \leq n$.

Consider the sequence $\{x_j\}_{j \in \mathbb{N}}$:

\hookrightarrow **Lemma 3.4.** $\{x_j\}$ cannot have a convergent subsequence.

Proof. Follows by $d(x_m, x_n) \geq \varepsilon \forall m, n$. ■

This contradicts our assumption that X sequentially compact, and we conclude X must be totally bounded. ■

⊗ **Example 3.3: Complete Metric Space Example: L^p norm**

Let $f \in C([a, b])$. We define the norm

$$\|f\|_p := \left(\int_a^b |f(x)|^p dx \right)^{\frac{1}{p}}.$$

As desired, $\|f\|_p \geq 0$; $\|f\|_p = 0 \iff f \equiv 0$; $\|c \cdot f\|_p = c \cdot \|f\|_p$.

Hölder's and Minkowski's inequalities for functions also hold; for $\frac{1}{p} + \frac{1}{q} = 1$, $1 \leq p, q \leq \infty$,

$$\int |fg| \leq \|f\|_p \cdot \|g\|_q; \quad \|f + g\|_p \leq \|f\|_p + \|g\|_p,$$

respectively.

We similarly have the L^∞ norm, namely, for a function $f : [a, b] \rightarrow \mathbb{R}$,

$$\|f\|_\infty = \sup_{x \in [a, b]} |f(x)|,$$

which obeys all the necessary properties as well.

Let $f_n \rightarrow f$ in $C([a, b])$, wrt $\|\cdot\|_\infty$, where $\{f_n\}_{n \in \mathbb{N}}$ a sequence of functions. Namely, we say that

$$\forall \varepsilon > 0, \exists N \in \mathbb{N} : \forall n \geq N, \sup_{x \in [a, b]} |f_n(x) - f(x)| < \varepsilon.$$

If this holds, we say that f_n *uniformly converges*.

We say that $f_n(x) \rightarrow f(x)$ *pointwise* on $[a, b]$ if $\forall x \in [a, b], f_n(x) \rightarrow f(x)$. Note that uniform convergence implies pointwise convergence, but not the converse.

↔ **Theorem 3.5**

Suppose $f_n(x)$ continuous, and $f_n(x) \rightarrow f(x)$ uniformly on $[a, b]$. Then, $f(x)$ also continuous on $[a, b]$.

Proof. Fix $\varepsilon > 0$, $x_0 \in [a, b]$. We have that $\exists N : n \geq N, |f_n(x) - f(x)| < \frac{\varepsilon}{3}, \forall x \in [a, b]$.

Let $n \geq N$. $f_n(x)$ continuous at x_0 , hence $\exists \delta(x_0) > 0 : |y - x_0| \implies |f_n(y) - f_n(x_0)| < \frac{\varepsilon}{3}$. We have

$$\begin{aligned} |f(x_0) - f(y)| &\leq |f(x_0) - f_n(x_0)| + |f_n(x_0) - f_n(y)| + |f_n(y) - f(y)| \\ &\leq \frac{\varepsilon}{3} + \frac{\varepsilon}{3} + \frac{\varepsilon}{3} = \varepsilon, \end{aligned}$$

completing the proof. ■

Remark 3.8. *This does not hold with pointwise convergence.*

Remark 3.9. We will prove later that $C([a, b])$ is complete for $\|f\|_\infty$, but not for arbitrary $\|f\|_p$, $1 \leq p < +\infty$. To “complete” $C([a, b])$ for $p \neq \infty$, we will need to consider measurable functions and redefine our notion of integration.

↪ Lecture 11; Last Updated: Thu Feb 8 09:51:13 EST 2024

4 DERIVATIVES

4.1 Introduction

↪ Definition 4.1: Differentiable

We say $f(x)$ differentiable at c if $\exists \lim_{x \rightarrow c} \frac{f(x)-f(c)}{x-c}$. If so, we denote the limit $f'(c)$.

Remark 4.1. For x close to c , then $f(x) \approx f(c) + f'(c)(x - c)$; this is a linear approximation of f at c .

⊗ Example 4.1: Weierstrass

$f(x) = \sum_{n=1}^{\infty} \frac{\cos(3^n x)}{2^n}$ is continuous in \mathbb{R} , but nowhere differentiable.

↪ Definition 4.2

The derivative, dx , is a linear map $C^1([a, b]) \rightarrow C^0([a, b])$.

4.2 Chain Rule

Remark 4.2. See [Analysis I notes](#) as well.

↪ Theorem 4.1: Caratheodory's Theorem

Let $f : I \rightarrow \mathbb{R}$, $c \in I$. f is differentiable at $x = c$ iff $\exists \varphi(x) : I \rightarrow \mathbb{R}$ s.t. φ continuous at c and $f(x) - f(c) = \varphi(x)(x - c)$.⁵

Proof. If $f'(c)$ exists, let

$$\varphi(x) = \begin{cases} \frac{f(x)-f(c)}{x-c} & x \neq c \\ f'(c) & x = c. \end{cases}$$

which is well defined. Moreover, for $x \neq c$, $\varphi(x)(x - c) = \frac{f(x)-f(c)}{x-c}(x - c) = f(x) - f(c)$ as desired; the case for $x = c$ is clear. Continuity at c :

$$\lim_{x \rightarrow c} \varphi(x) = \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} = f'(c) = \varphi(c).$$

⁵If not stated otherwise, sets named I or J are intervals.

Conversely, suppose such a φ exists. Then, by continuity,

$$\exists \varphi(c) = \lim_{x \rightarrow c} \varphi(x) = \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c}$$

which gives directly that f differentiable at c . ■

↪ Theorem 4.2: Chain Rule

Let $f : J \rightarrow \mathbb{R}, g : I \rightarrow \mathbb{R}, f(J) \subseteq I$. If $f(x)$ differentiable at c and $g(y)$ is differentiable at $y = f(c)$, then $g \circ f(x)$ is also differentiable at c , and

$$(g \circ f)'(c) = g'(f(c))f'(c).$$

Proof. Using [Caratheodory's Theorem](#), $\exists \varphi : f(x) - f(c) = \varphi(x)(x - c)$ with $\varphi(c) = f'(c)$. Let $d = f(c)$, then similarly $\exists \psi : g(y) - g(d) = \psi(y)(y - d)$ with $\psi(d) = g'(d)$, with φ, ψ continuous at c, d resp. Then,

$$g(f(x)) - g(f(c)) = \psi(f(x))(f(x) - f(c)) = (\psi \circ f)(x) \cdot (\varphi(x)(x - c))$$

$\psi \circ f$ is continuous at c , as a composition of continuous functions (ψ, ϕ continuous by construction, f differentiable and thus continuous). It follows, then, that

$$\lim_{x \rightarrow c} \frac{g(f(x)) - g(f(c))}{x - c} = \lim_{x \rightarrow c} (\psi \circ f)(x) \cdot \varphi(x) = \psi(f(c))\varphi(c) = g'(f(c)) \cdot f'(c),$$

by construction. ■

4.3 Critical Points

↪ Definition 4.3

$f : I \rightarrow \mathbb{R}$ has a max/min c if $\exists J \subseteq I : x \in J$ s.t. $\max_{x \in J} f(x) / \min_{x \in J} f(x) = f(c)$.

↪ Theorem 4.3: Rolle's

Let $f : [a, b] \rightarrow \mathbb{R}$ continuous. Suppose $f'(x)$ exists for all $x \in (a, b)$ and $f(a) = f(b) = 0$. Then, $\exists c \in (a, b) : f'(c) = 0$.

Remark 4.3. A "complex-version" of Rolle's:

↪ Theorem 4.4: Gauss-Lucas

Let $P(z)$ be a complex-valued polynomial. Then, the roots of $P'(z)$ lie inside the convex hull of roots of $P(z)$, where a convex hull is the smallest polygon with vertices at the roots of $P(z)$.

↪ **Definition 4.4**

Consider $P(z) = z^n - 1$ for some $n \in \mathbb{N}$. If z a root, we can show that $(|z|)^n = 1$, hence all roots lie on the unit circle in the complex plane at multiples of the same angle. This gives us a regular n -gon in the complex plane. We then have that $P'(z) = nz^{n-1}$, with has root $z = 0$, which clearly lies within the n -gon hull.

↪ **Theorem 4.5: Mean Value**

Let f be continuous on $[a, b]$ and differentiable on (a, b) . Then, $\exists c \in (a, b)$ s.t. $f(b) - f(a) = f'(c)(b - a)$.

Proof. Let $\varphi(x) = f(x) - f(a) = \frac{f(b)-f(a)}{(b-a)}(x - a)$, where $\varphi(a) = 0 = \varphi(b)$. By Rolle's theorem, $\exists c \in (a, b) : \varphi'(c) = 0 = f'(c) - \frac{f(b)-f(a)}{(b-a)}$, as desired. ■

↪ Lecture 12; Last Updated: Tue Apr 9 22:31:27 EDT 2024

4.4 Aside: Continued Fractions

We have that, for any $x \in \mathbb{R}$, $x = \lfloor x \rfloor + \{x\}$, with $\{x\} \in (0, 1)$; $\lfloor x \rfloor$ and $\{x\}$ are the integral and fractional parts of x respectively.

Fix $x \in \mathbb{R}$, assuming $x \neq 0$.

Let $x_1 := \frac{1}{\{x\}}$. We can write

$$x = \lfloor x \rfloor + \frac{1}{x_1}.$$

If $\{x_1\} \neq 0$, let $x_2 := \frac{1}{\{x_1\}}$ and write

$$x = \lfloor x \rfloor + \frac{1}{\lfloor x_1 \rfloor + \frac{1}{x_2}}.$$

Continuing in this manner, this process stops if $\{x_i\} = 0$ for some i ; if $x \in \mathbb{Q}$, this process will stop, else, it will continue infinitely. For instance, the Golden Ratio $x = \frac{\sqrt{5} \pm 1}{2}$ has continued fraction expansion

$$x = \frac{1}{1 + \frac{1}{1 + \frac{1}{\dots}}}$$

More succinctly, we can denote $a_0 := \lfloor x \rfloor$ and $a_i = \lfloor x_i \rfloor, i \geq 1$, and write

$$x = a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \ddots}}}$$

We notate, accordingly, $x := (a_1, a_2, a_3, \dots)$; in this case, the Golden Ratio can be notated $(1, 1, 1, \dots)$.

We denote $\frac{p_n}{q_n}$ as the n th continued fraction of a given x . It turns out that this is the best possible rational approximation for $x \notin \mathbb{Q}$.

4.5 Back To Derivatives

↪ Theorem 4.6

$f : I \rightarrow \mathbb{R}$, differentiable. f is increasing (resp decreasing) iff $f'(x) \geq 0 \forall x \in I$ (resp $f'(x) \leq 0 \forall x \in I$).

↪ Proposition 4.1

Let f continuous on $I = [a, b]$. Let $a < c < b$ and suppose f differentiable on (a, c) and (c, b) . Suppose $f'(x) \geq 0$ on $(c - \delta, c)$ and $f'(x) \leq 0$ on $(c, c + \delta)$ for some $\delta > 0$. Then, f has local max at $x = c$.

↪ Lemma 4.1

Let $I \subseteq \mathbb{R}$, and assume $f : I \rightarrow \mathbb{R}$ is differentiable at $x = c \in I$.

1. If $f'(c) > 0$, then $\exists \delta > 0 : f(x) > f(c) \forall x \in I, x \in (c, c + \delta)$.
2. (Reverse statement for $f'(c) < 0$)

↪ Theorem 4.7: Darboux

Suppose f differentiable on $I := [a, b]$ and $f'(a) < k < f'(b)$. Then, $\exists c \in (a, b)$ such that $f'(c) = k$.

↪ Lecture 13; Last Updated: Thu Feb 15 09:49:55 EST 2024

4.6 L'Hopital's Rules

↪ Proposition 4.2

Suppose $f(x), g(x) : [a, b] \rightarrow \mathbb{R}$ with $f(a) = g(a) = 0$, and $g(x) \neq 0 \forall a < x < b$. Suppose f, g are differentiable at $x = a$ and $g'(a) \neq 0$. Then, $\lim_{x \rightarrow a^+} \frac{f(x)}{g(x)}$ exists, and moreover, it is equal to $\frac{f'(a)}{g'(a)}$.

Proof.

$$\lim_{x \rightarrow a^+} \left(\frac{f(x) - f(a)}{x - a} \right) / \left(\frac{g(x) - g(a)}{x - a} \right) = \lim_{x \rightarrow a^+} \frac{f(x) - f(a)}{x - a} \cdot \frac{x - a}{g(x) - g(a)} = \lim_{x \rightarrow a^+} \frac{f(x) - f(a)}{g(x) - g(a)}$$

but the original line is simply $\frac{f'(a)}{g'(a)}$. ■

⊗ Example 4.2

$$\lim_{x \rightarrow 0} \frac{\sin x}{x} = \frac{\cos(0)}{1} = 1.$$

↪ Theorem 4.8: Cauchy Mean Value

Let $f(x), g(x) : [a, b] \rightarrow \mathbb{R}$ where f, g continuous on $[a, b]$ and differentiable on (a, b) . Assuming $g'(x) \neq 0, \forall x \in (a, b)$, then $\exists c \in (a, b)$ such that

$$\frac{f(b) - f(a)}{g(b) - g(a)} = \frac{f'(c)}{g'(c)}.$$

↪ Proposition 4.3: More General L'Hopital

let $-\infty \leq a < b \leq +\infty$ and f, g differentiable on (a, b) . Suppose $\lim_{x \rightarrow a^+} f(x) = 0 = \lim_{x \rightarrow a^+} g(x)$.

1. If $\exists L := \lim_{x \rightarrow a^+} \frac{f'(x)}{g'(x)}$ where L some real number, then $\lim_{x \rightarrow a^+} \frac{f(x)}{g(x)} = L$ as well.
2. If $\exists L := \lim_{x \rightarrow a^+} \frac{f'(x)}{g'(x)}$ where $L = +\infty$ or $-\infty$, then $\lim_{x \rightarrow a^+} \frac{f(x)}{g(x)} = L$ as well.

↪ Proposition 4.4

Let $-\infty \leq a < b \leq +\infty, f, g$ differentiable on (a, b) and $g'(x) \neq 0 \forall x \in (a, b)$. Suppose $\lim_{x \rightarrow a^+} g(x) = \pm\infty$.

1. If $\lim_{x \rightarrow a^+} \frac{f'(x)}{g'(x)} =: L$ exists and is some finite real number, then $\lim_{x \rightarrow a^+} \frac{f(x)}{g(x)} = L$ as well.
2. If $\lim_{x \rightarrow a^+} \frac{f'(x)}{g'(x)} =: L$ exists and is $\pm\infty$, then $\lim_{x \rightarrow a^+} \frac{f(x)}{g(x)} = L$ as well.

4.7 Taylor's Theorem

↪ Theorem 4.9: Taylor's Theorem

Let $I = [a, b] \subseteq \mathbb{R}, f : I \rightarrow \mathbb{R}, f \in C^n(I)$ and suppose $f^{(n+1)}(x)$ exists on (a, b) . Let $x_0 \in [a, b]$. Then, for any $x \in [a, b], \exists c$ between x, x_0 such that

$$f(x) = f(x_0) + f'(x_0)(x - x_0) + \frac{f''(x_0)}{2}(x - x_0)^2 + \dots + \frac{f^{(n)}(x_0)}{n!}(x - x_0)^n + \frac{f^{(n+1)}(c)}{(n+1)!}(x - x_0)^{n+1}$$

↪ Theorem 4.10: Relative Extrema

⁶Let $I \subseteq \mathbb{R}$ be an open interval, $x_0 \in I$, and $n \geq 2$. Suppose $f', f'', \dots, f^{(n)}$ are continuous in a neighborhood of x_0 , and $f'(x_0) = f''(x_0) = \dots = f^{(n-1)}(x_0) = 0$ and $f^{(n)}(x_0) \neq 0$. Then:

1. if n is even and $f^{(n)}(x_0) > 0$, then f has a local minimum at x_0 ;
2. if n is even and $f^{(n)}(x_0) < 0$, then f has a local maximum at x_0 ;
3. if n is odd, then f has neither a local minimum nor maximum at x_0 .

Proof. If $n := 2m$ -even and $f^{(2m)}(x_0) > 0$, then $f^{(n)}(c) > 0$ so $f(x) - f(x_0) = f^{(2m)}(c)(x - x_0) > 0$. ■

4.8 Convex Sets

↪ Definition 4.5: Convex Set

$A \subseteq V$ -vector space over \mathbb{R} is *convex* if for any $x, y \in A$ and any $0 \leq t \leq 1$, $t \cdot x + (1 - t) \cdot y \in A$.

↪ Definition 4.6: Convex Function

Let $f : I \rightarrow \mathbb{R}$. f is *convex* if $\forall x_1, x_2 \in I$ and $0 \leq t \leq 1$,

$$f((1 - t)x_1 + tx_2) \leq (1 - t)f(x_1) + tf(x_2).$$

↪ Lecture 15; Last Updated: Thu Feb 22 21:53:23 EST 2024

5 RIEMANN INTEGRAL

5.1 Introduction

↪ Definition 5.1: Partitions

A *partition* is a division of an interval (a, b) , denoted

$$\mathcal{P} = \{a = x_0, x_1, \dots, x_{n-1}, x_n = b\}.$$

We define $\text{diam}(\mathcal{P}) := \max_n |x_i - x_{i-1}|$.

A *marked partition*, denoted $\dot{\mathcal{P}}$, is one in which, for each interval we choose some $t_i \in (x_i, x_{i+1}]$.

⁶Bartle-Sherbert, Theorem 6.4.4

↪ **Definition 5.2: Riemann Sum**

We denote

$$S(f, \dot{\mathcal{P}}) = \sum_{i=1}^n f(t_i)(x_i - x_{i-1}).$$

↪ **Definition 5.3: Riemann Integrable**

A function f is *Riemann Integrable* on $[a, b]$ if $S(f, \dot{\mathcal{P}}) \rightarrow L$ as $\text{diam}(\dot{\mathcal{P}}) \rightarrow 0$ for any choice of $t_i \in [x_i, x_{i+1}]$.

That is, $\forall \varepsilon > 0, \exists \delta : \text{if } \text{diam}(\mathcal{P}) < \delta, \text{ then for any choice of } t_i \in [x_i, x_{i+1}] \text{ we have } |L - S(f, \dot{\mathcal{P}})| < \varepsilon.$

↪ **Proposition 5.1**

1. If L exists, it is unique.
2. The integral is linear in $f(x)$; if $\int_a^b f(x) dx$ and $\int_a^b g(x) dx$ exist, then $\int_a^b (c_1 f + c_2 g) dx = c_1 \int_a^b f dx + c_2 \int_a^b g dx$.
3. If $f \leq g$ are Riemann integrable on $[a, b]$, then $\int_a^b f(x) dx \leq \int_a^b g(x) dx$.

↪ **Proposition 5.2**

If $f(x)$ integrable on $[a, b]$, the $f(x)$ is bounded on $[a, b]$.

Proof. Suppose $\int_a^b f$ exists. Let $\varepsilon > 0$, and δ such that if $\text{diam}(\dot{\mathcal{P}}) < \delta$ then $|L - S(f, \dot{\mathcal{P}})| < \varepsilon$. Let $\varepsilon = 1$. Then, $S(f, \dot{\mathcal{P}}) \leq |L| + 1$.

Let $Q = \{a = x_0, \dots, x_n = b\}$ be a partition of $[a, b]$ such that $\text{diam}(Q) < \delta$. Suppose towards a contradiction that f is not bounded on $[a, b]$. Then, f is unbounded on at least one interval $[x_i, x_{i+1}]$, say, on $[x_k, x_{k+1}]$. Let $t_i = x_i$ for $i \neq k$ and choose $t_k \in [x_k, x_{k+1}]$ such that $|f(t_k)|(x_{k+1} - x_k) > |L| + 1 + |\sum_{i \neq k} f(t_i)(x_{i+1} - x_i)|$ (which we can do by assumption of f being unbounded).

By assumption, $|S(f, \dot{Q})| \leq |L| + 1$, but we have that

$$S(f, \dot{Q}) = \underbrace{\sum_{i \neq k} f(t_i)(x_{i+1} - x_i)}_{:=N} + |f(t_k)|(x_{k+1} - x_k) > 2N + |L| + 1,$$

contradiction. ■

5.2 Cauchy Criterion

↪ **Proposition 5.3: Cauchy Criterion for Integrability**

$f \in \mathcal{R}[a, b] \iff \forall \varepsilon > 0, \exists \delta > 0 : \text{if } \dot{P} \text{ and } \dot{Q} \text{ are tagged partitions of } [a, b] \text{ s.t. } \text{diam } \dot{P} < \delta \text{ and } \text{diam } \dot{Q} < \delta, \text{ then } |S(f, \dot{P}) - S(f, \dot{Q})| < \varepsilon.^7$

5.3 Squeeze Theorem

↪ **Theorem 5.1**

Let $f : [a, b] \rightarrow \mathbb{R}$. Then $\int_a^b f$ exists $\iff \forall \varepsilon > 0, \exists \alpha_\varepsilon(x), \omega_\varepsilon(x) \in \mathcal{R}[a, b], \alpha_\varepsilon \leq f \leq \omega_\varepsilon$, and

$$\int_a^b (\omega_\varepsilon - \alpha_\varepsilon) < \varepsilon$$

Proof. If $f \in \mathcal{R}[a, b]$ then take $\alpha_\varepsilon = f = \omega_\varepsilon$.

Conversely, let $\varepsilon > 0$. Since $\alpha_\varepsilon, \omega_\varepsilon \in \mathcal{R}[a, b]$, then, $\exists \delta > 0$ such that for any tagged partition with $\text{diam } \dot{P} < \delta$, then $|S(\alpha_\varepsilon, \dot{P}) - \int_a^b \alpha_\varepsilon| < \varepsilon$ and $|S(\omega_\varepsilon, \dot{P}) - \int_a^b \omega_\varepsilon| < \varepsilon$, thus

$$\int_a^b \alpha_\varepsilon - \varepsilon < S(\alpha_\varepsilon, \dot{P}) \leq S(f, \dot{P}) \leq S(\omega_\varepsilon, \dot{P}) < \int_a^b \omega_\varepsilon + \varepsilon.$$

Let \dot{Q} be any other tagged partition with $\text{diam } \dot{Q} < \delta$; then, the same inequality holds ie $\int_a^b \alpha_\varepsilon - \varepsilon < S(f, \dot{Q}) < \int_a^b \omega_\varepsilon + \varepsilon$. Subtracting one from the other, we see that

$$|S(f, \dot{P}) - S(f, \dot{Q})| < \int_a^b \omega_\varepsilon - \int_a^b \alpha_\varepsilon + 2\varepsilon < 3\varepsilon,$$

and thus $f \in \mathcal{R}[a, b]$ by Cauchy Criterion. ■

↪ Lecture 16; Last Updated: Wed Apr 10 08:39:04 EDT 2024

↪ **Lemma 5.1: BS-7.2.4**

Let $J \subseteq [a, b]$ an interval with endpoints $c < d$. If

$$\varphi_J(x) := \begin{cases} 1 & x \in J \\ 0 & x \notin J \end{cases}.$$

Then, $\varphi_J \in \mathcal{R}[a, b]$ and $\int_a^b \varphi_J = d - c$.

⁷Note that $\mathcal{R}[a, b]$ is the set of all real-valued functions integrable on the interval $[a, b]$.

↪ **Theorem 5.2**

Let $\varphi : [a, b] \rightarrow \mathbb{R}$, $\varphi \in \mathcal{R}[a, b]$; that is, step functions are integrable.

↪ **Theorem 5.3**

f continuous on $[a, b]$ implies $f \in \mathcal{R}[a, b]$.

Proof. (Sketch) f uniform continuous; use this to construct step functions that “bound” f from above and below. Apply the squeeze theorem. ■

↪ Lecture 17; Last Updated: Thu Mar 28 14:33:47 EDT 2024

↪ **Theorem 5.4: BS-7.2.7**

Monotone functions on $[a, b]$ are integrable.

Proof. We show only for increasing. Let $f : [a, b] \rightarrow \mathbb{R}$ be monotone increasing. If f constant, then it is a step function and we are done.

Otherwise, $f(b) - f(a) > 0$. Let $\varepsilon > 0$ and $q \in \mathbb{N}$ such that $h := \frac{f(b)-f(a)}{q} < \frac{\varepsilon}{b-a}$, effectively subdividing the y -axis into q equal-sized parts. Then, let

$$y_i := f(a) + ih, \quad 0 \leq i \leq q,$$

and take

$$A_k := f^{-1}([y_{k+1}, y_k]) = \begin{cases} \emptyset \\ \{x\} \\ I_i \end{cases} .$$

We disregard each $A_k : A_k = \emptyset$, and adjoin the isolated points $\{x\}$ to the I_i 's, and hence have a partition $\cup_k A_k = [a, b]$. Letting $\alpha(x) = y_{k-1}$ and $\omega(x) = y_k$ for $x \in A_k$, then $\alpha(x) \leq f(x) \leq \omega(x) \forall x \in [a, b]$ (effectively, we are created a “series of squeezes”). Then,

$$\int_a^b \omega(x) - \alpha(x) dx = \sum_{k=1}^q (y_k - y_{k-1})(x_k - x_{k-1}) = h(b - a) < \varepsilon,$$

and the proof is completed by applying the squeeze theorem. ■

↪ **Theorem 5.5: Additivity; BS-7.2.8**

Let $f : [a, b] \rightarrow \mathbb{R}$ and $a < c < b$. Then, $f \in \mathcal{R}[a, b] \iff f \in \mathcal{R}[a, c]$ and $f \in \mathcal{R}[c, b]$. Moreover, $\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$.

Proof. See book. Remark that this holds for finite summations of integrals as such by induction. ■

5.4 Fundamental Theorem of Calculus

↪ Definition 5.4

Call $F(x)$ a *primitive* of $f(x)$ if F differential and $F'(x) = f(x)$.

↪ Theorem 5.6: Fundamental Theorem of Calculus

Let $F, f : [a, b] \rightarrow \mathbb{R}$ and $E \subseteq [a, b]$ a finite set s.t.

1. F continuous on $[a, b]$
2. $F'(x) = f(x) \forall x \in [a, b] \setminus E$; ie they agree for all but finitely many points
3. $f \in \mathcal{R}[a, b]$

Then, $\int_a^b f(x) = F(b) - F(a)$.

Proof. (Sketch) Remark first that it suffices to prove for $E := \{a, b\}$; using additivity, we can subdivide any other such E into such subsets of 1 or 2 elements.

Fix $\varepsilon > 0$ and take $\delta > 0$ such that for any \dot{P} of $[a, b]$ s.t. $\text{diam } \dot{P} < \delta$, $\left| S(f, \dot{P}) - \int_a^b f(x) \right| < \varepsilon$. Applying the mean value theorem to F on each $[x_{i-1}, x_i]$ of \dot{P} :

$$\begin{aligned} F(x_i) - F(x_{i-1}) &= F'(u_i)(x_i - x_{i-1}), \quad u_i \in [x_{i-1}, x_i] \\ &= f(u_i)(x_i - x_{i-1}) \end{aligned}$$

Hence, summing over each of these,

$$\begin{aligned} F(x_1) - \cancel{F(x_0)} + \overset{F(a)}{\cancel{F(x_2)}} - F(x_1) + \cdots + \cancel{F(x_n)} - \overset{F(b)}{\cancel{F(x_{n-1})}} &= f(u_1)(x_1 - a) + \cdots + f(u_n)(x_n - x_{n-1}) \\ \implies F(b) - F(a) &= \sum_{i=1}^n f(u_i)(x_i - x_{i-1}) =: S(f, \dot{P}_1) \end{aligned}$$

by construction, $\text{diam}(\dot{P}_1) < \delta$ since the only change we have made from \dot{P} is the tags, hence $\left| S(f, \dot{P}_1) - \int_a^b f(x) \right| < \varepsilon$. Thus,

$$\begin{aligned} \left| S(f, \dot{P}_1) - \int_a^b f(x) \right| &= \left| F(b) - F(a) - \int_a^b f(x) \right| < \varepsilon \\ \implies F(b) - F(a) &= \int_a^b f(x) \quad \text{as } \varepsilon \rightarrow 0. \end{aligned}$$

■

5.5 Upper and Lower Riemann Sums

↪ Definition 5.5: Upper/Lower Riemann Sums

For a partition P ,

- $\bar{S}(f, P) := \sum_{i=1}^n (\sup_{t \in [x_{i-1}, x_i]} f(t)) \cdot (x_i - x_{i-1})$
- $\underline{S}(f, P) := \sum_{i=1}^n (\inf_{t \in [x_{i-1}, x_i]} f(t)) \cdot (x_i - x_{i-1})$

↪ Proposition 5.4

For any tagged partition \dot{P} ,

$$\underline{S}(f, P) \leq S(f, \dot{P}) \leq \bar{S}(f, P).$$

Moreover, $f \in \mathcal{R}[a, b]$ if $\forall \varepsilon > 0, \exists \delta > 0$ s.t. $\text{diam}(P) < \delta \implies \left| \bar{S}(f, P) - \underline{S}(f, P) \right| < \varepsilon$.

Proof. (Sketch) Remark that this is a similar idea to saying that $\inf = \sup \implies$ limit exists. ■

↪ Proposition 5.5

Let P_1, P_2 be partitions of $[a, b]$, and let P_3 be the *common refinement* of P_1, P_2 . Then

$$\underline{S}(f, P_i) \leq \underline{S}(f, P_3) \leq \bar{S}(f, P_3) \leq \bar{S}(f, P_i), \quad i = 1, 2,$$

that is, the finer refinement always gives a better approximation.

5.6 Indefinite Integral

↪ Definition 5.6

For $f \in \mathcal{R}[a, b]$ and any $z \in [a, b]$, define

$$F(z) := \int_a^z f(x) dx.$$

↪ Theorem 5.7

$F(z)$ continuous on $[a, b]$.

Proof. $f \in \mathcal{R}[a, b] \implies f$ bounded $\implies \exists M$ s.t. $|f(x)| \leq M \forall x \in [a, b]$, so (assuming $z < w$),

$$|F(z) - F(w)| = \left| \int_a^z f(x) dx - \int_a^w f(x) dx \right| = \left| \int_z^w f(x) dx \right| \leq M \cdot |z - w|,$$

so taking $w \rightarrow z$, $|F(z) - F(w)| \rightarrow 0$. ■

↪ **Theorem 5.8: Another Fundamental Theorem of Calculus**

Let $f \in \mathcal{R}[a, b]$, f -continuous at $c \in [a, b]$. Then $F(z)$ differentiable at c and $F'(c) = f(c)$.

↪ **Corollary 5.1**

If $f(x)$ continuous on $[a, b]$ $F'(x) = f(x) \forall x \in [a, b]$.

↪ **Theorem 5.9: Substitution/Change of Variables**

Let $J := [\alpha, \beta]$, $\varphi : J \rightarrow \mathbb{R}$, $\varphi \in C^1([a, b])$. Suppose $\varphi(J) \subseteq I \subseteq \mathbb{R}$, and let $f : I \rightarrow \mathbb{R}$ be continuous on I . Then,

$$\int_{\varphi(\alpha)}^{\varphi(\beta)} f(x) dx = \int_{\alpha}^{\beta} f(\varphi(t)) \cdot \varphi'(t) dt .$$

Proof. Left as a (homework) exercise; make use of the chain rule! ■

⊗ **Example 5.1**

Compute $\int_1^4 \frac{\sin(\sqrt{t})}{\sqrt{t}} dt$ using the previous theorem.

5.7 Lebesgue Integrability Criterion

↪ **Definition 5.7: Lebesgue Measure 0**

$A \subseteq \mathbb{R}$ has *Lebesgue measure 0* iff $\forall \varepsilon > 0$, A can be covered by a countable union of intervals $J_k := [a_k, b_k]$ such that $\sum_{k=1}^{\infty} |J_k| \leq \varepsilon$. We also call such an A a null set.

For some set $S \subseteq \mathbb{R}$ and statement P , we say “ P holds for almost every $x \in S$ ” if $\{x \in S : P \text{ false}\}$ has Lebesgue measure 0.

⊗ **Example 5.2**

1. Any countable set is a null set.
2. The Cantor set is a null set.

↪ **Theorem 5.10: Lebesgue Integrability Criterion**

Let $f : [a, b] \rightarrow \mathbb{R}$ be a bounded function. Then

$$\begin{aligned} f \in \mathcal{R}[a, b] &\iff f \text{ - continuous for almost every } x \in [a, b] \\ &\iff \{z \in [a, b] : f \text{ discontinuous}\} \text{ has Lebesgue measure } 0. \end{aligned}$$

Remark 5.1. The proof is rather involved, but is in the appendix of Bartle. Its important to remark that this is a necessary and sufficient condition.

⊗ **Example 5.3**

1. Let $f : [0, 1] \rightarrow \mathbb{R}, f(x) := \begin{cases} 1 & x \in \mathbb{Q} \\ 0 & x \notin \mathbb{Q} \end{cases}$. f discontinuous everywhere, so $f \notin \mathcal{R}[a, b]$.

2. Let $f(x) := \begin{cases} \frac{1}{b} & x = \frac{a}{b} \in \mathbb{Q} \text{ s.t. } (a, b) = 1 \\ 0 & x \notin \mathbb{Q} \end{cases}$. One can show that f continuous on $x \in \mathbb{R} \setminus \mathbb{Q}$ and only discontinuous on \mathbb{Q} . But this is a countable set so certainly has Lebesgue measure 0 and so $f \in \mathcal{R}[0, 1]$.

↪ Lecture 19; Last Updated: Tue Apr 9 14:45:17 EDT 2024

↪ **Theorem 5.11: Composition**

$f \in \mathcal{R}[a, b], f([a, b]) \subseteq [c, d], \varphi : [c, d] \rightarrow \mathbb{R}$ continuous, then $\varphi \circ f \in \mathcal{R}[a, b]$.

Proof.

$$\{x \text{ s.t. } \varphi \circ f \text{ discontinuous at } x\} \subseteq \{x : f \text{ discontinuous at } x\}$$

since φ continuous. The RHS has Lebesgue measure 0, and thus so does the LHS, hence the proof. ■

↪ **Theorem 5.12: Product Theorem**

$f, g \in \mathcal{R}[a, b] \implies f \cdot g \in \mathcal{R}[a, b]$.

Proof. $f \cdot g = \frac{1}{4} [(f + g)^2 - (f - g)^2]$. $f \pm g \in \mathcal{R}[a, b]$ and so so is $(f \pm g)^2$ by taking $\varphi(x) := x^2$ as in the previous theorem. It follows that $f \cdot g \in \mathcal{R}[a, b]$. ■

5.8 Integration by Parts

↪ **Theorem 5.13**

Let F, G be differentiable on $[a, b]$, with $f := F', g := G'$. Suppose $f, g \in \mathcal{R}[a, b]$, then

$$\int_a^b f(x)G(x) dx = F(x)G(x)|_a^b - \int_a^b F(x)g(x) dx.$$

Proof. Remark that $(FG)' = F'G + FG' = fG + Fg$, so on the one hand

$$\int_a^b (FG)' dx = \int_a^b (fG + Fg) dx = \int_a^b fG dx + \int_a^b Fg dx,$$

but on the other hand, by the fundamental theorem of calculus,

$$\int_a^b (FG)' dx = [F \cdot G](a) - [F \cdot G](b) = F(x)G(x)|_a^b,$$

and so

$$\begin{aligned} F(x)G(x)|_a^b &= \int_a^b fG dx + \int_a^b Fg dx \\ \implies \int_a^b f(x)G(x) dx &= F(x)G(x)|_a^b - \int_a^b F(x)g(x) dx, \end{aligned}$$

and hence the result. ■

↪ **Theorem 5.14: Taylor's Theorem, Remainder's Version**

Suppose $f', f'', \dots, f^{(n)}$ exist on $[a, b]$ and $f^{(n+1)} \in \mathcal{R}[a, b]$.⁸ Then,

$$f(b) = f(a) + \frac{f'(a)}{1!}(b-a) + \frac{f''(a)}{2!} + \dots + \frac{f^{(n)}(a)}{n!}(b-a)^n + R_n,$$

with $R_n := \frac{1}{n!} \int_a^b f^{(n+1)}(t)(b-t)^n dt$.

Proof. See Bartle; makes use of integration by parts. ■

6 FUNCTION SEQUENCES, SERIES

6.1 Pointwise and Uniform Convergence

⁸Remark that this is a weaker condition than continuity as was used in our previous statement of Taylor's theorem.

↪ **Definition 6.1: Pointwise vs Uniform Convergence**

We say a sequence of functions $f_n \rightarrow f$ *pointwise* on a set E if $\forall x \in E, f_n(x) \rightarrow f(x)$ as $n \rightarrow \infty$.

On the other hand, $f_n \rightarrow f$ *uniformly* on E if $\forall \varepsilon > 0, \exists N \in \mathbb{N}$ such that $\forall n \geq N$ and $x \in E$, $|f_n(x) - f(x)| < \varepsilon$.

Remark 6.1. Notice that uniformly implies pointwise convergence.

⊗ **Example 6.1**

Let $f_n := \begin{cases} 2nx & 0 \leq x \leq \frac{1}{2n} \\ 0 & x > \frac{1}{2n} \end{cases}$. Show that $f_n \rightarrow 0$ pointwise but not uniformly (hint: $f_n(\frac{1}{2n}) = 1 \forall n$).

↪ **Theorem 6.1**

The space of functions $C([a, b])$ equipped with the sup norm is complete.

Proof. Proven in tutorials. ■

↪ **Theorem 6.2: Interchange of Limits**

Let $J \subseteq \mathbb{R}$ be a bounded interval such that $\exists x_0 \in J : f_n(x_0) \rightarrow f(x_0)$. Suppose $f'_n(x) \rightarrow g(x)$ uniformly $\forall x \in J$. Then, $\exists f : f_n(x) \rightarrow f(x)$ uniformly on J , $f(x)$ differentiable on J , and $f'(x) = g(x) \forall x \in J$.

Proof. This is a rather painful proof; one needs to make use of the “multiple epsilons” from each given continuity/convergence/differentiability statement. ■

↪ Lecture 20; Last Updated: Wed Apr 10 13:31:53 EDT 2024

↪ **Theorem 6.3**

Let $f_n \in \mathcal{R}[a, b], f_n \rightarrow f$ uniformly on $[a, b]$. Then, $f \in \mathcal{R}[a, b]$ and $\int_a^b f_n(x) dx \rightarrow \int_a^b f(x) dx$.

↪ **Theorem 6.4: Bounded Convergence Theorem**

$f_n \in \mathcal{R}[a, b], f_n \rightarrow f \in \mathcal{R}[a, b]$, not necessarily uniformly. Suppose $\exists B > 0$ s.t. $|f_n(x)| \leq B \forall x \in [a, b]$. Then, $\int_a^b f_n \rightarrow \int_a^b f$ as $n \rightarrow \infty$.

↪ **Theorem 6.5: Dini's Theorem/Monotone Convergence**

$f_n \in C([a, b]), f_n(x)$ monotone (as a sequence). Suppose $f_n \rightarrow f \in C([a, b])$. Then, $f_n \rightarrow f$ uniformly on $[a, b]$.

↪ Lecture 21; Last Updated: Thu Mar 28 11:58:26 EDT 2024

6.2 Series

↪ Definition 6.2: Absolute Convergence

Let $\{x_j\} \in X$ where X a normed vector space (say, \mathbb{R}). We say

$$\sum_{j=1}^{\infty} x_j \text{ converges absolutely} \iff \sum_{j=1}^{\infty} \|x_j\| < +\infty.$$

↪ Theorem 6.6

Any rearrangement of absolutely convergent series given the same sum.

↪ Definition 6.3: Conditional Convergence

$\sum_{j=1}^{\infty} \vec{x}^{(j)}$ conditionally convergent if $\sum_{j=1}^{\infty} x^{(j)}$ converges (ie each component converges) but $\sum_{j=1}^{\infty} \|\vec{x}^{(j)}\| = \infty$.

↪ Theorem 6.7

If $\sum_{i=1}^{\infty} a_i \in \mathbb{R}$ conditionally convergent, you can change the order of summation such that $\forall x \in \mathbb{R}$, $\exists \sigma$ -permutation such that $\sum_{i=1}^{\infty} a_{\pi(i)} = x$.

Proof. (Sketch) Separate a_i into positive, negative parts. Since conditionally convergent, $\sum_{a_j > 0} a_j = +\infty$ and $\sum_{a_j < 0} a_j = -\infty$. Add positive a_i 's until the partial sum $\geq x$, then add negative a_i 's until the partial sum $\leq x$, and repeat. The final rearrangement will converge as desired. ■

↪ Lecture 22; Last Updated: Thu Mar 28 12:14:46 EDT 2024

↪ Theorem 6.8

Suppose $\sum_{i=1}^{\infty} \vec{v}_i, \vec{v}_i \in \mathbb{R}^n$, converges, but $\sum_{i=1}^{\infty} \|\vec{v}_i\| = +\infty$. Then, the set of rearranged sums $\sum_{i=1}^{\infty} \vec{v}_{\sigma(i)}$ for each $\sigma : \mathbb{N} \leftrightarrow \mathbb{N}$ permutation form an *affine subspace* of \mathbb{R}^n .

6.3 Tests for Absolute Convergence

↪ Proposition 6.1

Let x_n, y_n be sequences and $r := \lim_{n \rightarrow \infty} \left| \frac{x_n}{y_n} \right|$.

1. If $r \neq 0$, $\sum_{n=1}^{\infty} x_n$ converges absolutely iff $\sum_{n=1}^{\infty} y_n$ converges absolutely. In addition, if $0 < r_1 := \liminf \left| \frac{x_n}{y_n} \right| \leq \limsup \left| \frac{x_n}{y_n} \right| =: r_2 < +\infty$, this still holds.
2. If $r = 0$, and if $\sum y_n$ converges absolutely, so does $\sum x_n$.

↪ **Proposition 6.2: Root Test**

If there $\exists r < 1$ such that $|x_n|^{1/n} \leq r$ for sufficiently large $n \geq K$, then $\sum_{n=K}^{\infty} |x_n| \leq \sum_{n=K}^{\infty} r^{-n}$ converges.

If $|x_n|^{1/n} \geq n$ for $n \geq K$, $\sum x_n$ does not converge absolutely.

↪ **Proposition 6.3: Ratio Test**

Let $x_n \neq 0$. If $\exists 0 < r < 1$, $\left| \frac{x_{n+1}}{x_n} \right| \leq r$ for sufficiently large n , $\sum x_n$ absolutely convergent.

If $\left| \frac{x_{n+1}}{x_n} \right| \geq 1$ for $n \geq K$, $\sum x_n$ diverges.

↪ **Proposition 6.4: Integral Test**

Let $f(x) \geq 0$ be non-increasing/non-decreasing function of $x \geq 1$. Then $\sum_{k=1}^{\infty} f(k)$ converges $\iff \lim_{k \rightarrow \infty} \int_1^k f(x) dx$ finite.

↪ **Proposition 6.5: Raube's Test**

Let $x_n \neq 0$.

1. Suppose $\exists a > 1$ s.t. $\left| \frac{x_{n+1}}{x_n} \right| \leq 1 - \frac{1}{n}$, $n \geq K$. Then $\sum x_n$ converges absolutely.
2. If $\exists a \leq 1$ s.t. $\left| \frac{x_{n+1}}{x_n} \right| \geq 1 - \frac{1}{n}$, $n \geq K$. Then $\sum x_n$ does not converge absolutely.

↪ **Corollary 6.1**

Let $a := \lim n \left(1 - \left| \frac{x_{n+1}}{x_n} \right| \right)$, if such a limit exists. Then, if $a > 1$, $\sum x_n$ converges absolutely, and if $a < 1$, $\sum x_n$ does not.

6.4 Tests for Non-Absolute Convergence

↪ **Proposition 6.6: Alternating Series**

If $x_n > 0$, $x_{n+1} \leq x_n$, $\lim_{n \rightarrow \infty} x_n = 0 \implies \sum (-1)^n x_n$ converges.

↪ **Lemma 6.1: Abel's Lemma**

Let $x_n, y_n \in \mathbb{R}$. Let $s_0 := 0$, $s_n := \sum_{k=1}^n y_k$. Then, for $m > n$,

$$\sum_{k=n+1}^m x_k y_k = x_m s_m - x_{n+1} s_{n+1} + \sum_{k=n+1}^m (x_k - x_{k+1}) s_k$$

↪ **Theorem 6.9: Dirichlet's Test**

Suppose x_n decreasing and $\lim_{n \rightarrow \infty} x_n = 0$. If $s_n := y_1 + \dots + y_n$ bounded, then $\sum_{n=1}^{\infty} x_n y_n$ converges.

Proof. Fix $B > 0$ such that $|s_n| \leq B$. x_n decreasing so $x_k - x_{k+1} \geq 0$. By Abel's lemma,

$$\begin{aligned} \left| \sum_{k=n+1}^m x_k y_k \right| &\leq |x_m s_m - x_{n+1} s_n| + \left| \sum_{k=n+1}^{m-1} s_k (x_k - x_{k+1}) \right| \\ &\leq x_m B + x_{n+1} B + \underbrace{\sum_{k=n+1}^{m-1} (x_k - x_{k+1}) B}_{\text{telescopes}} \\ &= 2x_{n+1} B \xrightarrow{n \rightarrow \infty} 0. \end{aligned}$$

Remark 6.2. *What is red and commutes? An abelian grape!*

⊗ **Example 6.2: Improving Convergence**

$y_j = (-1)^{j+1}$ does not converge, but $|s_n| \leq 1$, so taking $x_n = \frac{1}{n}$ gives $\sum_n x_n y_n$ finite.

This is an example of "improving convergence", ie making a nearly-convergent series convergence. Another example is by taking successive arithmetic means of a given sequence; ie let $a_1 = y_1, a_2 = \frac{y_1+y_2}{2}, \dots, a_n = \frac{\sum_{i=1}^n y_i}{n}$. Then, in this case again $y_n = (-1)^{n+1}$, which does not converge, has $a_n \rightarrow 0$.

Moreover, if $y_n \rightarrow A$, then $a_n \rightarrow A$ as well (converse does not hold in general, as above).

↪ **Theorem 6.10: Abel's Test**

Let x_n -convergent and monotone, and suppose $\sum_n y_n$ converges. Then $\sum_n x_n y_n$ also converges.

6.5 Series of Functions

↪ **Definition 6.4: Convergence**

We say a series of functions $\sum_n f_n(x)$ absolutely convergent on E if $\sum_n |f_n(x)|$ converges for all $x \in E$.

We say the convergence $\sum_n f_n(x) \rightarrow g(x)$ is uniform if the convergence is uniform for any $x \in E$; that is, $\forall \varepsilon > 0, \exists N \in \mathbb{N}$ such that $\forall n \geq N$ and $x \in E, |g(x) - \sum_n f_n(x)| < \varepsilon$.

↪ **Proposition 6.7**

Suppose for $f_n : E := [a, b] \rightarrow \mathbb{R}$, $\sum_n f_n(x) \rightarrow g(x)$ uniform for $x \in E$, and $f_n \in \mathcal{R}[a, b]$. Then

$$\int_a^b g(x) dx = \sum_{n=1}^{\infty} \int_a^b f_n(x) dx .$$

That is, the integral of the limit is equal to the limit of the integral.

↪ **Proposition 6.8**

Let $f_n : [a, b] \rightarrow \mathbb{R}$ where $f'_n \exists$ on $[a, b]$. Suppose $\sum_n f_n(x)$ converges for some $x \in [a, b]$ and $\sum_n f'_n(x)$ converges uniformly on $[a, b]$. Then there exists some $g : [a, b] \rightarrow \mathbb{R}$ such that $\sum_n f_n \rightarrow g$ uniformly on $[a, b]$, g differentiable on $[a, b]$, and $g'(x) = \sum_n f'_n(x)$. That is, the derivative of the limit equals the limit of the derivatives.

↪ Lecture 24; Last Updated: Wed Apr 10 15:09:45 EDT 2024

↪ **Theorem 6.11: Cauchy Criterion**

$f_n(x) : D \subseteq \mathbb{R} \rightarrow \mathbb{R}$ converges uniformly on D iff $\forall \varepsilon > 0 \exists N$ s.t. $\forall m, n \geq N, \sum_{i=n+1}^m f_i(x) < \varepsilon \forall x \in D$.

↪ **Proposition 6.9: Weierstrass M-Test**

If $|f_n(x)| \leq M_n \forall x \in D \subseteq \mathbb{R}$ and $\sum_n M_n < +\infty$, then $\sum_n f_n(x)$ converges uniformly on D .

Proof. Suffices to look at the tail: $\sum_{j=n+1}^m |f_n(x)| \leq \sum_{j=n+1}^m M_j$. ■

6.6 Power Series

↪ **Definition 6.5: Power Series**

A function, series of the form

$$f(x) = \sum_{n=0}^{\infty} a_n(x - c)^n \quad \circledast$$

is said to be a power series centered at $c \in \mathbb{R}$.

↪ **Definition 6.6: Radius of Convergence**

For a_n as in \circledast , let $\rho := \limsup_{n \rightarrow \infty} \sqrt[n]{|a_n|}$. Then, $R := \frac{1}{\rho}$ the radius of convergence of f (taking $R = 0$ if $\rho = \infty$, $R = \infty$ if $\rho = 0$).

↔ **Theorem 6.12: Cauchy-Hadamard**

Let R be the radius of convergence of \otimes . Then, $\sum_n a_n(x - c)^n$ converges if $|x - c| < R$, and diverges if $|x - c| > R$. If precisely equal, either case could happen (and needs to be treated individually).

Proof. Directly apply the root test; assume $c = 0$. Then $|a_n x^n|^{1/n} = |a_n|^{1/n} |x|$. If $|x| < \rho$, then $\limsup |a_n|^{1/n} |x| < \rho^{1/\rho} = 1$ so $\forall \varepsilon > 0, \exists N$ s.t. $\forall n \geq N, |a_n|^{1/n} |x| < 1 - \varepsilon \implies |a_n| |x|^n < (1 - \varepsilon)^n$. It follows that $\sum_{j \geq N} |a_n x^n| < \sum_{j \geq N} (1 - \varepsilon)^j$. But this RHS is a geometric series with $r < 1$ so thus converges. The converse follows similarly (well, backwards). ■

⊗ Example 6.3

1. $1 + x + x^2 + \dots$ converges absolutely for $|x| < 1$.
2. $1 + x + \frac{x^2}{2} + \frac{x^3}{3} + \dots$ converges for $-1 \leq x < 1$.
3. $\sum_n \frac{x^n}{n^k}$ with $k \geq 2$ converges for $-1 \leq x \leq 1$ (check the $x = 1$ case by comparison test, then the $x = -1$ test follows by alternating series test.)

↔ **Theorem 6.13**

Let J be a closed and bounded interval strictly contained in the interval of convergence of \otimes . Then $f(x)$ converges uniformly in J .

↔ Lecture 25; Last Updated: Tue Apr 9 15:14:47 EDT 2024

Remark 6.3. *In-class review. Good luck!*

↔ Lecture 26; Last Updated: Tue Apr 9 15:16:14 EDT 2024

~

7 APPENDIX

7.1 Notes from Tutorials

↪ Theorem 7.1

Let (X, d) be a compact metric space.⁹ Let $C(X) := \{f : X \rightarrow \mathbb{R} : f \text{ continuous}\}$ be a vector space. Take the uniform norm $\|f\| := \sup_{x \in X} |f(x)|$ on $C(X)$. Then, $(C(X), \|\cdot\|)$ is complete.¹⁰

Proof. Denote the “canonical norm” $\rho(f, g) := \|f - g\|$.

Let $(f_n) \in C(X)$ be a Cauchy sequence. Then, $\forall \varepsilon > 0, \exists N \in \mathbb{N} : \forall m, n \geq N, \rho(f_n, f_m) < \varepsilon$.

Fix $x \in X$, noting that

$$|f_n(x) - f_m(x)| \leq \sup_{y \in X} |f_n(y) - f_m(y)| = \rho(f_n, f_m) < \varepsilon. \quad *^1$$

Define, for this fixed x , a sequence in $\mathbb{R} \{f_n(x)\}_{n \in \mathbb{N}}$. By $*^1$, we have that this sequence is Cauchy in \mathbb{R} , but as \mathbb{R} complete, $f_n(x)$ hence converges, to some limit we call $f(x) := \lim_{n \rightarrow \infty} f_n(x)$. Note that x is still fixed at this point; these are but real numbers we are working with here.

Now, as x was completely arbitrary, we can repeat this process for all of X , and define a function $f : X \rightarrow \mathbb{R}$ where $f(x) := \lim_{n \rightarrow \infty} f_n(x)$.

For a fixed x , we have that $f_m(x) \rightarrow f(x)$ as $m \rightarrow \infty$. This implies:

$$\begin{aligned} 0 \leq \lim_{m \rightarrow \infty} |f_n(x) - f_m(x)| &\leq \lim_{m \rightarrow \infty} \varepsilon = \varepsilon \\ &\implies |f_n(x) - f(x)| \leq \varepsilon \forall n \geq N \\ &\implies \rho(f_n, f) = \sup_{x \in X} |f_n(x) - f(x)| \leq \varepsilon \implies f_n \rightarrow f \end{aligned}$$

It remains to show that $f \in C(X)$. Let $c \in X$ and $\varepsilon > 0$, and the corresponding $N \in \mathbb{N} : \rho(f_n, f) < \frac{\varepsilon}{3} \forall n \geq N$. By construction, $f_N \in C(X)$, and is thus continuous at c . This gives that $\exists \delta > 0 : |f_N(x) - f_N(c)| < \frac{\varepsilon}{3}$ whenever $d(x, c) < \delta$.¹¹

Hence, if $d(x, c) < \delta$, we have

$$\begin{aligned} |f(x) - f(c)| &\leq |f(x) - f_N(x)| + |f_N(x) - f_N(c)| + |f_N(c) - f(c)| \\ &\leq \rho(f, f_N) + \frac{\varepsilon}{3} + \frac{\varepsilon}{3} \\ &< \frac{\varepsilon}{3} + \frac{\varepsilon}{3} + \frac{\varepsilon}{3} = \varepsilon, \end{aligned}$$

hence f continuous at c , which was completely arbitrary, and thus $f \in C(X)$. ■

¹⁰In this proof, the compactness is necessary for the norm to be well-defined.

¹⁰In this way, this becomes a Banach Space: a complete, normed vector space.

¹¹Be careful here, there are three different metrics going on; ρ from the vector space, d from the underlying metric space, and $|\cdot|$ from \mathbb{R} .

↪ **Theorem 7.2**

Let (X, d) -complete. Let $\{F_n\}$ be a decreasing family of non-empty closed sets with $\lim_{n \rightarrow \infty} \text{diam}(F_n) = 0$. Then, $\exists z : \bigcap_{n \in \mathbb{N}} F_n = \{z\}$.

↪ **Theorem 7.3**

Let (X, d) -complete, and $f : X \rightarrow X$ an “expanding map”, such that $d(x, y) \leq d(f(x), f(y)) \forall x, y \in X$. Then, f is a surjective isometry, ie, $f(X) = X$ and $d(f(x), f(y)) = d(x, y) \forall x, y \in X$.

↪ **Lemma 7.1**

Differentiable \implies Continuous.

Proof. Let $f : I \rightarrow \mathbb{R}$, and $c \in I$ arbitrary. Notice that $\forall x \neq c \in I, f(x) - f(c) = (x - c) \frac{f(x) - f(c)}{x - c}$. Hence,

$$\begin{aligned} \lim_{x \rightarrow c} (f(x) - f(c)) &= \lim_{x \rightarrow c} (x - c) \frac{f(x) - f(c)}{x - c} \\ &= \lim_{x \rightarrow c} (x - c) \cdot \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} \\ &= 0 \cdot f'(c) = 0 \\ &\implies \lim_{x \rightarrow c} f(x) = f(c), \end{aligned}$$

hence f continuous, noting that the splitting of the limits is valid as both are defined. ■

⊗ **Example 7.1**

$$\text{Let } f : \mathbb{R} \rightarrow \mathbb{R}, f(x) := \begin{cases} x^2 & x \in \mathbb{Q} \\ 0 & x \notin \mathbb{Q} \end{cases}$$

Claim: f discontinuous at all $x \neq 0$.

Proof. Let $x \neq 0 \in \mathbb{R}$. By density of $\mathbb{Q} \subseteq \mathbb{R}$, there exist sequences $(r_n) \in \mathbb{Q}$ s.t. $r_n \rightarrow x$ and $(z_n) \in \mathbb{R} \setminus \mathbb{Q}$ s.t. $z_n \rightarrow x$. Then:

$$\lim_{n \rightarrow \infty} f(r_n) = \lim_{n \rightarrow \infty} r_n^2 = x^2$$

$$\lim_{n \rightarrow \infty} f(z_n) = \lim_{n \rightarrow \infty} 0,$$

hence f discontinuous by the sequential criterion at $x \neq 0$. ■

Claim: $f'(0) = 0$.

Proof. Let $\varepsilon > 0$ and $\delta = \varepsilon$. Notice that $f(x) \leq x^2 \forall x$. Then, we have that $\forall |x| < \delta$,

$$\begin{aligned} \left| \frac{f(x) - f(0)}{x - 0} - 0 \right| &= \left| \frac{f(x)}{x} \right| \\ &\leq \left| \frac{x^2}{x} \right| = |x| < \delta = \varepsilon \end{aligned}$$

↪ **Definition 7.1**

Let $f : I \rightarrow \mathbb{R}$. A point $c \in I$ is a local max (resp min) if $\exists \delta > 0$ s.t. $f(x) \leq f(c)$ (resp $f(x) \geq f(c)$) $\forall x \in (c - \delta, c + \delta) \cap I$.

↪ **Lemma 7.2**

Let $f : I \rightarrow \mathbb{R}$ be differentiable at $c \in I^\circ$. If c a local extrema of f , then $f'(c) = 0$.

Proof. Assume wlog that c a local max; if a local min, take $\tilde{f} := -f$ and continue.

Since I° open, $\exists \delta_1 > 0 : (c - \delta_1, c + \delta_1) \subseteq I^\circ \subseteq I$. We also have that $\exists \delta_2 > 0 : f(x) \leq f(c) \forall x \in (c - \delta_2, c + \delta_2) \cap I$, by c an extrema.

Let $\delta := \min\{\delta_1, \delta_2\}$. Then, we have both $(c - \delta, c + \delta) \subseteq I$ and $f(x) \leq f(c) \forall x \in (c - \delta, c + \delta)$.

Since $f'(c)$ exists, $\lim_{x \rightarrow c^+} \frac{f(x) - f(c)}{x - c} = \lim_{x \rightarrow c^-} \frac{f(x) - f(c)}{x - c}$. But we have from the property of being a maximum

that

$$\lim_{x \rightarrow c^+} \frac{f(x) - f(c)}{x - c} \geq 0, \quad \lim_{x \rightarrow c^-} \frac{f(x) - f(c)}{x - c} \leq 0,$$

hence, as these two limits must agree, they must equal 0 and thus $f'(c) = 0$. ■

7.2 Miscellaneous

⊗ Example 7.2: Rudin, Chapter 7: Differentiability

1. Let f be defined $\forall x \in \mathbb{R}$, and suppose that $|f(x) - f(y)| \leq (x - y)^2, \forall x, y \in \mathbb{R}$. Prove that f is constant.¹²

Proof. Let $x > y \in \mathbb{R}$. Then, as $|x - y| = x - y$, we have

$$\begin{aligned} |f(x) - f(y)| \leq (x - y)^2 &\implies \left| \frac{f(x) - f(y)}{x - y} \right| \leq x - y = |x - y| \rightarrow 0 \text{ as } y \rightarrow x \\ &\implies \left| \frac{f(x) - f(y)}{x - y} \right| \rightarrow 0 \end{aligned}$$

This implies, then, that $f'(x)$ is defined $\forall x \in \mathbb{R}$, and moreover, that $f'(x) = 0 \forall x \in \mathbb{R}$. We conclude, then, that $f(x)$ constant $\forall x \in \mathbb{R}$. ■

2. Suppose $f'(x) > 0$ in (a, b) . Prove that f is strictly increasing in (a, b) , and let g be its inverse function. Prove that g is differentiable, and that

$$g'(f(x)) = \frac{1}{f'(x)} \quad (a < x < b).$$

Proof. Fix $x > y \in (a, b)$. Then, by the mean value theorem, $\exists z \in (x, y) : f'(z) = \frac{f(x) - f(y)}{x - y}$. Since $f'(z) > 0$, it follows that

$$\frac{f(x) - f(y)}{x - y} > 0 \implies f(x) - f(y) > x - y > 0 \implies f(x) > f(y),$$

hence, f increasing, as $x > y$ arbitrary.

Let now $g := f^{-1}$. ■

¹²Note that this means that f Hölder continuous with constant $\alpha = 2$. Indeed, Hölder continuous functions with $\alpha > 1$ are always constant by a similar proof. For $0 < \alpha \leq 1$, we have the inclusion continuously differentiable \implies Lipschitz \implies α -Hölder \implies uniformly continuous \implies continuous.

7.3 Class Midterm Solutions

↪ Question 7.1

Let X be a topological space, and let $f, g : X \rightarrow \mathbb{R}$ be two continuous functions. Show that the set $\{x \in X : f(x) > g(x)\}$ is an open subset of X .

Proof. Let $A := \{x \in X : f(x) > g(x)\}$. Letting $\varphi(x) := f(x) - g(x) = (f - g)(x)$, then remark that $A \equiv \{x \in X : \varphi(x) > 0\}$, and since differences of continuous functions are continuous, φ continuous. Letting $B := (0, \infty) \subseteq \mathbb{R}$, then, we have that $A = \varphi^{-1}(B)$. But B an open set, and the inverse images of open sets by continuous functions are open, hence A open. ■

↪ Question 7.2

- List three equivalent properties (definitions) of compact sets in metric spaces; you don't need to prove anything.
- Is the unit ball¹³ in the space ℓ^2 of infinite sequences compact? Prove or disprove. You may use any of the properties from (a).

Proof. (a) Every open cover admits a finite subcover \iff sequentially compact \iff complete and totally bounded.

- Denote the closed unit ball centered at $(0, 0, \dots)$ in ℓ^2 , $B := \{x \in \ell^2 : d_2^2(0, x) = \sum_{i=1}^{\infty} |x_i|^2 \leq 1\}$. Consider the sequence of "unit sequences"

$$\{e^n\}_{n \in \mathbb{N}} \in B, \quad e_i^n := \delta_{in}.$$

Then, for any $i \neq j$, $d_2(e^n, e^m) = \sqrt{2} > 1$. It follows that, although $e^n \in B$ for any n , there cannot exist a subsequence of x^n that converges within B (verify why this is!). Thus, B cannot be sequentially compact and thus not compact. ■

¹³Jakobson said in class this is supposed to be a closed ball.

↪ Question 7.3

- (a) Define a complete metric space.
- (b) State (without proof) the contraction mapping theorem.
- (c) Let $f : (0, 1) \rightarrow (0, 1)$ be defined by $f(x) = x/2$. Is f a contraction?
- (d) Does f have a fixed point in the open interval $I = (0, 1)$? Does that contradict the contraction mapping theorem?

Proof. (a) A complete metric space is a metric space in which every Cauchy sequence converges within that space.

(b) Let (X, d) be a complete metric space, and let $f : X \rightarrow X$ be a contraction mapping, ie for any $x, y \in X$, $d(f(x), f(y)) \leq c \cdot d(x, y)$ for some $c \in (0, 1)$. Then, the contraction mapping states that f has a unique fixed point $z \in X$, ie $f(z) = z$ and $\lim_{n \rightarrow \infty} f^{(n)}(x) = z$ for any x .

(c) For any $x, y \in (0, 1)$, we have

$$d(f(x), f(y)) = |f(x) - f(y)| = \left| \frac{x - y}{2} \right| = \frac{1}{2} |x - y| = c \cdot d(x, y),$$

so f indeed a contraction mapping with $c := \frac{1}{2}$.

(d) We have that for any $x \in I$, $f^{(n)}(x) = \frac{x}{2^n}$ so x a fixed point iff $\frac{x}{2^n} = \frac{x}{2^{n-1}}$ for some n , which is only possible if $x = 0$, but $0 \notin I$, so indeed f has no fixed point in I . This is not a contradiction to the contraction mapping theorem since $I := (0, 1)$ not complete (indeed, $\frac{1}{n} \in I \forall n$ but $\frac{1}{n} \rightarrow 0 \notin I$).

■

↪ **Question 7.4**

Let $x = (x_1, x_2, \dots)$ and $y = (y_1, y_2, \dots)$ be infinite real sequences satisfying $\|x\|_2 \leq 2$ and $\|y\|_2 \leq 3$, where $\|\cdots\|_2$ the ℓ^2 norm.

(a) State Holder's inequality and Minkowski inequality for sequences.

(b) Give an upper bound for $x \cdot y = \sum_i x_i y_i$, and for $\|x + y\|$.

Proof. (a) Holder's inequality: for p, q Holder conjugates and $x \in \ell^p, y \in \ell^q$ we have

$$\left| \sum_{i=1}^{\infty} x_i y_i \right| \leq \|x\|_p \|y\|_q.$$

Minkowski inequality: for $x, y \in \ell^p$,

$$\|x + y\|_p \leq \|x\|_p + \|y\|_p.$$

(b) For x, y as given; by Holders, $x \cdot y \leq \|x\|_p \|y\|_q = 2 \cdot 3 = 6$, and by Minkowski's, $\|x + y\| \leq \|x\| + \|y\| = 2 + 3 = 5$, so 6, 5 are upper bounds for $x \cdot y, \|x + y\|$ respectively.

■

↪ Question 7.5

(a) State (without proof) Taylor's theorem.

(b) Let $f \in C^4([0, 2])$, and let $f'(1) = f''(1) = f'''(1) = 0$ while $f^{(4)}(1) = 2$. Use (a) to show that $f(x)$ has a local extremum at $x = 1$, and determine its type.

Proof. (a) Let $I := [a, b] \subseteq \mathbb{R}$ and let $f : I \rightarrow \mathbb{R}$ such that $f \in C^n(I)$, and $f^{(n+1)}(x)$ exists on (a, b) . Then, for $x_0 \in [a, b]$, there exists some $c \in (\min(x, x_0), \max(x, x_0))$ such that

$$f(x) = f(x_0) + f'(x_0)(x - x_0) + \cdots + \frac{f^{(n)}(x_0)}{n!}(x - x_0)^n + \frac{f^{(n+1)}(c)}{(n+1)!}(x - x_0)^{n+1}.$$

(b) By Taylor's, for any $x \in [0, 2]$, there exists some c between x and 1 such that

$$\begin{aligned} f(x) &= f(1) + \underbrace{f'(1)(\cdots) + f''(1)(\cdots) + f'''(1)(\cdots)}_{=0} + \frac{f^{(4)}(c)}{4!}(x - 1)^4 \\ &= f(1) + \frac{f^{(4)}(c)}{4!}(x - 1)^4 \\ \implies f(x) - f(1) &\geq \frac{f^{(4)}(c)}{4!}(x - 1)^4 \forall x \in [0, 2] \end{aligned}$$

By continuity of $f^{(4)}$, there exists some neighborhood V of $x_0 = 1$ such that $f^{(4)}(c)$ has the same sign of $f^{(4)}(1)$. So, for any $x \in V$, $\frac{f^{(4)}(c)}{4!} \geq 0$, since $\frac{f^{(4)}(1)}{4!} = \frac{2}{4!} \geq 0$. Thus, since $(x - 1)^4 \geq 0$, we have that for such x in V ,

$$f(x) - f(1) \geq 0 \implies f(x) \geq f(1).$$

Hence, we have a neighborhood of 1 such that for all x in the neighborhood $f(x) \geq f(1)$. It follows that 1 a local minimum of f . ■

8 LIST OF THEOREMS

↪ Definition 1.1 (Metric Space)	3
↪ Definition 1.2 (Open Metric Space)	3
↪ Definition 1.3 (Normed Space)	3
↪ Proposition 1.1	3
↪ Proposition 1.2	4
↪ Definition 1.4 (Convex Set)	4
↪ Definition 1.5 (ℓ_p)	5
↪ Proposition 1.3	6
↪ Definition 2.1 (Topological space)	7
↪ Definition 2.2 (Closed sets)	7
↪ Definition 2.3 (Equivalence of Metrics)	8
↪ Definition 2.4 (Interior, Boundary of a Topological Set)	8
↪ Definition 2.5 (Closure)	8
↪ Proposition 2.1 (Properties of $\text{Int}(A)$)	9
↪ Proposition 2.2 (Properties of \overline{A})	9
↪ Proposition 2.3	9
↪ Definition 2.6 (Basis for a Topology)	9
↪ Proposition 2.4	9
↪ Proposition 2.5	10
↪ Definition 2.7	10
↪ Proposition 2.6 (Consequences of Subspace Topologies)	10
↪ Proposition 2.7	10
↪ Lemma 2.1	11
↪ Lemma 2.2	11
↪ Definition 2.8 (Continuous Function)	11
↪ Proposition 2.8	11
↪ Proposition 2.9	11
↪ Theorem 2.1 (Continuity of Composition)	12
↪ Proposition 2.10	12
↪ Definition 2.9 (Finite Product Spaces)	12
↪ Definition 2.10 (Cylinder Set)	12
↪ Definition 2.11 (Projection)	13
↪ Definition 2.12 (Coordinate Function)	13
↪ Proposition 2.11	13
↪ Definition 2.13 (Product Topology/Cylinder Sets for ∞ Products)	14
↪ Proposition 2.12	14
↪ Proposition 2.13	15
↪ Question 2.1	15
↪ Definition 2.14 (Compact)	16
↪ Proposition 2.14	16
↪ Theorem 2.2 (On Compactness)	17
↪ Proposition 2.15	17
↪ Proposition 2.16	17
↪ Proposition 2.17	18
↪ Definition 2.15 (Connected)	18
↪ Theorem 2.3	18

↪ Lemma 2.3	19
↪ Theorem 2.4 (“Intermediate Value Theorem”)	19
↪ Theorem 2.5	19
↪ Theorem 2.6	20
↪ Definition 2.16 (Path Connected)	20
↪ Theorem 2.7 (Path connected \implies connected)	20
↪ Proposition 2.18	20
↪ Definition 2.17 (Connected Component)	21
↪ Definition 2.18 (Path Component)	21
↪ Proposition 2.19	21
↪ Lemma 2.4	21
↪ Lemma 2.5	21
↪ Lemma 2.6	21
↪ Proposition 2.20	22
↪ Proposition 2.21	22
↪ Definition 2.19 (An Explicit Definition)	22
↪ Definition 2.20 (Complement Definition)	22
↪ Definition 3.1 (Hölder Conjugates)	23
↪ Proposition 3.1 (Hölder’s Inequality)	23
↪ Lemma 3.1	24
↪ Proposition 3.2 (Minkowski Inequality)	26
↪ Theorem 3.1	27
↪ Definition 3.2 (Complete Metric Space)	27
↪ Proposition 3.3	27
↪ Proposition 3.4	28
↪ Proposition 3.5	28
↪ Proposition 3.6	28
↪ Definition 3.3 (Contraction Mapping)	29
↪ Theorem 3.2 (Contraction Mapping Theorem)	29
↪ Theorem 3.3 (ℓ_p complete)	30
↪ Definition 3.4 (Totally Bounded)	32
↪ Theorem 3.4	32
↪ Lemma 3.1 (Cantor Intersection Theorem)	32
↪ Definition 3.5 (Finite Intersection Property)	33
↪ Lemma 3.2 (Finite Intersection Formulation of Compactness)	33
↪ Lemma 3.3	33
↪ Lemma 3.4	33
↪ Theorem 3.5	34
↪ Definition 4.1 (Differentiable)	35
↪ Definition 4.2	35
↪ Theorem 4.1 (Caratheodory’s Theorem)	35
↪ Theorem 4.2 (Chain Rule)	36
↪ Definition 4.3	36
↪ Theorem 4.3 (Rolle’s)	36
↪ Theorem 4.4 (Gauss-Lucas)	36
↪ Definition 4.4	37
↪ Theorem 4.5 (Mean Value)	37
↪ Theorem 4.6	38
↪ Proposition 4.1	38
↪ Lemma 4.1	38

↪ Theorem 4.7 (Darboux)	38
↪ Proposition 4.2	38
↪ Theorem 4.8 (Cauchy Mean Value)	39
↪ Proposition 4.3 (More General L'Hopital)	39
↪ Proposition 4.4	39
↪ Theorem 4.9 (Taylor's Theorem)	39
↪ Theorem 4.10 (Relative Extrema)	40
↪ Definition 4.5 (Convex Set)	40
↪ Definition 4.6 (Convex Function)	40
↪ Definition 5.1 (Partitions)	40
↪ Definition 5.2 (Riemann Sum)	41
↪ Definition 5.3 (Riemann Integrable)	41
↪ Proposition 5.1	41
↪ Proposition 5.2	41
↪ Proposition 5.3 (Cauchy Criterion for Integrability)	42
↪ Theorem 5.1	42
↪ Lemma 5.1 (BS-7.2.4)	42
↪ Theorem 5.2	43
↪ Theorem 5.3	43
↪ Theorem 5.4 (BS-7.2.7)	43
↪ Theorem 5.5 (Additivity; BS-7.2.8)	43
↪ Definition 5.4	44
↪ Theorem 5.6 (Fundamental Theorem of Calculus)	44
↪ Definition 5.5 (Upper/Lower Riemann Sums)	45
↪ Proposition 5.4	45
↪ Proposition 5.5	45
↪ Definition 5.6	45
↪ Theorem 5.7	45
↪ Theorem 5.8 (Another Fundamental Theorem of Calculus)	46
↪ Corollary 5.1	46
↪ Theorem 5.9 (Substitution/Change of Variables)	46
↪ Definition 5.7 (Lebesgue Measure 0)	46
↪ Theorem 5.10 (Lebesgue Integrability Criterion)	47
↪ Theorem 5.11 (Composition)	47
↪ Theorem 5.12 (Product Theorem)	47
↪ Theorem 5.13	48
↪ Theorem 5.14 (Taylor's Theorem, Remainder's Version)	48
↪ Definition 6.1 (Pointwise vs Uniform Convergence)	49
↪ Theorem 6.1	49
↪ Theorem 6.2 (Interchange of Limits)	49
↪ Theorem 6.3	49
↪ Theorem 6.4 (Bounded Convergence Theorem)	49
↪ Theorem 6.5 (Dimi's Theorem/Monotone Convergence)	49
↪ Definition 6.2 (Absolute Convergence)	50
↪ Theorem 6.6	50
↪ Definition 6.3 (Conditional Convergence)	50
↪ Theorem 6.7	50
↪ Theorem 6.8	50
↪ Proposition 6.1	50
↪ Proposition 6.2 (Root Test)	51

↪ Proposition 6.3 (Ratio Test)	51
↪ Proposition 6.4 (Integral Test)	51
↪ Proposition 6.5 (Raube's Test)	51
↪ Corollary 6.1	51
↪ Proposition 6.6 (Alternating Series)	51
↪ Lemma 6.1 (Abel's Lemma)	51
↪ Theorem 6.9 (Dirichlet's Test)	52
↪ Theorem 6.10 (Abel's Test)	52
↪ Definition 6.4 (Convergence)	52
↪ Proposition 6.7	53
↪ Proposition 6.8	53
↪ Theorem 6.11 (Cauchy Criterion)	53
↪ Proposition 6.9 (Weierstrass M-Test)	53
↪ Definition 6.5 (Power Series)	53
↪ Definition 6.6 (Radius of Convergence)	53
↪ Theorem 6.12 (Cauchy-Hadamard)	54
↪ Theorem 6.13	54
↪ Theorem 7.1	55
↪ Theorem 7.2	56
↪ Theorem 7.3	56
↪ Lemma 7.1	56
↪ Definition 7.1	57
↪ Lemma 7.2	57
↪ Question 7.1	59
↪ Question 7.2	59
↪ Question 7.3	60
↪ Question 7.4	61
↪ Question 7.5	62